

Groups acting by conjugation

Recall that a group G acts on itself by conjugation as follows:

$$g \cdot a = gag^{-1} \quad \forall g, a \in G.$$

This satisfies the axioms for a group action:

$$1 \cdot a = |a|^{-1} = a, \text{ and}$$

$$(gh) \cdot a = (gh)a(gh)^{-1} = ghah^{-1}g^{-1} = g \cdot (h \cdot a).$$

Def: Elements $a, b \in G$ are conjugate if $\exists g$ s.t. $b = gag^{-1}$, i.e. if they are in the same orbit under the conjugation action. The orbits in this case are called the conjugacy classes of G .

Ex: If $a \in Z(G)$, then a is the only element in its conjugacy class. If $a \notin Z(G)$, then there is some g s.t. $gag^{-1} \neq a$, so there are at least two elements in its conjugacy class.

Note: If G is nontrivial, the action of conjugation can't be transitive, since the conjugacy class of 1 is always just $\{1\}$.

Recall that if G acts on A , and $a \in A$, we showed that the number of elts in its orbit will be equal to $|G : G_a|$, i.e. the index of its stabilizer in G .

When G acts on itself by conjugation, and $h \in G$,
 $C_h = \{g \in G \mid ghg^{-1} = h\} = C_G(s).$

That is,

Prop: The number of conjugates of an element $s \in G$ is $|G : C_G(s)|$.

We know that the orbits of an action partition the set being acted on, so in particular, if we add up the # of elements in all the orbits, we get the following:

Thm: (The Class equation) Let G be a finite group, and g_1, g_2, \dots, g_r representatives of the distinct conjugacy classes not contained in the center of G . Then

$$|G| = |Z(G)| + \sum_{i=1}^r |G : C_G(g_i)|$$

Pf: Each orbit in the center has exactly one element. If the other orbits are K_1, \dots, K_r and g_i a representative from K_i , then $|K_i| = |G : C_G(g_i)|$.

Since the orbits partition G , summing up their cardinalities gives us the desired equation. \square

Ex: In D_8 , the center is $\{1, r^2\}$.

The centralizer of r contains $\langle r \rangle$, so it has order ≥ 4 . Thus, $|G : C_G(r)| \leq \frac{8}{4} = 2$. But $srs = r^3$, so its conjugacy class is $\{r, r^3\}$.

$C_G(s) = \{1, s, r^2, sr^2\}$, so $|G : C_G(s)| = 2$, and $r sr^{-1} = sr^2$, so its conj. class is $\{s, sr^2\}$.

Note that the two remaining elts are conjugate: $r(sr)r^{-1} = sr^3$, so $\{sr, sr^3\}$ is the final conjugacy class.

Note that all of the summands in the class group divide the order of the group. This helps us classify some finite groups.

Theorem: If p is prime, and G is a group of order p^α , some $\alpha \geq 1$, then G has nontrivial center.

Pf: let g_1, \dots, g_r be representatives from the conjugacy classes not contained in the center (if there are any).

Then for each g_i , its conjugacy class has at least 2 elements, so $1 < |G : C_G(g_i)|$, and Lagrange's Thm says that the index must divide p^α . Thus, for each g_i , $p \mid |G : C_G(g_i)|$.

The class equation says that

$$p^n = |Z(G)| + \sum_{i=1}^r |G : C_G(g_i)|$$

\uparrow
divisible by p

Thus $p \mid |Z(G)|$, so $Z(G)$ is not trivial. \square

Cor: If $|G| = p^2$ for some prime p , then G is abelian, and G is cyclic or $G \cong \mathbb{Z}_p \times \mathbb{Z}_p$.

Pf: $|Z(G)| = p$ or p^2 by the above. Thus $|G/Z(G)| = 1$ or p , so it's cyclic. Thus, by a HW problem, G is abelian.

The nontrivial elements of G have orders p or p^2 . If G has any element of order p^2 , then G is cyclic. Thus, assume all nontrivial elements have order p .

Let $x \in G$ s.t. $x \neq 1$. Then $|x| = p$, so we can find $y \in G - \langle x \rangle$.

Then $\langle x \rangle \times \langle y \rangle \cong \mathbb{Z}_p \times \mathbb{Z}_p$. Define $\varphi: \langle x \rangle \times \langle y \rangle \rightarrow G$ by $(x^a, y^b) \mapsto x^a y^b$. It is straightforward to check this is a homomorphism.

If $(x^a, y^b) \in \ker \varphi$, then $x^a y^b = 1 \Rightarrow x^a = y^{-b} \in \langle x \rangle \cap \langle y \rangle$. But $\langle x \rangle \cap \langle y \rangle \neq \langle x \rangle$, and the order divides p , so $a = b = 0$.

Thus, $\ker \varphi = 1$, so φ is injective. Both groups have the same order, so it must also be a bijection and thus an isomorphism. \square