

# Math 221: LINEAR ALGEBRA

## Chapter 4. Vector Geometry

### §4-2. Projections and Planes

Le Chen<sup>1</sup>

Emory University, 2020 Fall

(last updated on 10/26/2020)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.

The Dot Product and Angles

Projections

Planes

Cross Product

Shortest Distances

NOTE: Much of this chapter is what you would learn in Multivariable Calculus.

You might find it interesting/useful to read.

But I will only cover the material important to this course.



# The Dot Product and Angles

## Definition

Let  $\vec{u} = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix}$  be vectors in  $\mathbb{R}^3$ . The **dot product** of  $\vec{u}$  and  $\vec{v}$  is

$$\vec{u} \cdot \vec{v} = x_1x_2 + y_1y_2 + z_1z_2,$$

i.e.,  $\vec{u} \cdot \vec{v}$  is a **scalar**.

# The Dot Product and Angles

## Definition

Let  $\vec{u} = \begin{bmatrix} x_1 \\ y_1 \\ z_1 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix}$  be vectors in  $\mathbb{R}^3$ . The **dot product** of  $\vec{u}$  and  $\vec{v}$  is

$$\vec{u} \cdot \vec{v} = x_1x_2 + y_1y_2 + z_1z_2,$$

i.e.,  $\vec{u} \cdot \vec{v}$  is a **scalar**.

## Remark

Another way to think about the dot product is as the  $1 \times 1$  matrix

$$\vec{u}^T \vec{v} = \begin{bmatrix} x_1 & y_1 & z_1 \end{bmatrix} \begin{bmatrix} x_2 \\ y_2 \\ z_2 \end{bmatrix} = \begin{bmatrix} x_1x_2 + y_1y_2 + z_1z_2 \end{bmatrix}.$$

## Theorem ( Properties of the Dot Product )

Let  $\vec{u}, \vec{v}, \vec{w}$  be vectors in  $\mathbb{R}^3$  (or  $\mathbb{R}^2$ ) and let  $k \in \mathbb{R}$ .

## Theorem ( Properties of the Dot Product )

Let  $\vec{u}, \vec{v}, \vec{w}$  be vectors in  $\mathbb{R}^3$  (or  $\mathbb{R}^2$ ) and let  $k \in \mathbb{R}$ .

1.  $\vec{u} \cdot \vec{v}$  is a real number.



## Theorem ( Properties of the Dot Product )

Let  $\vec{u}, \vec{v}, \vec{w}$  be vectors in  $\mathbb{R}^3$  (or  $\mathbb{R}^2$ ) and let  $k \in \mathbb{R}$ .

1.  $\vec{u} \cdot \vec{v}$  is a real number.

2.  $\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$ .

(commutative property)

## Theorem ( Properties of the Dot Product )

Let  $\vec{u}, \vec{v}, \vec{w}$  be vectors in  $\mathbb{R}^3$  (or  $\mathbb{R}^2$ ) and let  $k \in \mathbb{R}$ .

1.  $\vec{u} \cdot \vec{v}$  is a real number.
2.  $\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$ . (commutative property)
3.  $\vec{u} \cdot \vec{0} = 0$ .

## Theorem ( Properties of the Dot Product )

Let  $\vec{u}, \vec{v}, \vec{w}$  be vectors in  $\mathbb{R}^3$  (or  $\mathbb{R}^2$ ) and let  $k \in \mathbb{R}$ .

1.  $\vec{u} \cdot \vec{v}$  is a real number.
2.  $\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$ . (commutative property)
3.  $\vec{u} \cdot \vec{0} = 0$ .
4.  $\vec{u} \cdot \vec{u} = ||\vec{u}||^2$ .

## Theorem ( Properties of the Dot Product )

Let  $\vec{u}, \vec{v}, \vec{w}$  be vectors in  $\mathbb{R}^3$  (or  $\mathbb{R}^2$ ) and let  $k \in \mathbb{R}$ .

1.  $\vec{u} \cdot \vec{v}$  is a real number.
2.  $\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$ . (commutative property)
3.  $\vec{u} \cdot \vec{0} = 0$ .
4.  $\vec{u} \cdot \vec{u} = ||\vec{u}||^2$ .
5.  $(k\vec{u}) \cdot \vec{v} = k(\vec{u} \cdot \vec{v}) = \vec{u} \cdot (k\vec{v})$ . (associative property)

## Theorem ( Properties of the Dot Product )

Let  $\vec{u}, \vec{v}, \vec{w}$  be vectors in  $\mathbb{R}^3$  (or  $\mathbb{R}^2$ ) and let  $k \in \mathbb{R}$ .

1.  $\vec{u} \cdot \vec{v}$  is a real number.

2.  $\vec{u} \cdot \vec{v} = \vec{v} \cdot \vec{u}$ . (commutative property)

3.  $\vec{u} \cdot \vec{0} = 0$ .


4.  $\vec{u} \cdot \vec{u} = ||\vec{u}||^2$ .

5.  $(k\vec{u}) \cdot \vec{v} = k(\vec{u} \cdot \vec{v}) = \vec{u} \cdot (k\vec{v})$ . (associative property)


6.  $\vec{u} \cdot (\vec{v} + \vec{w}) = \vec{u} \cdot \vec{v} + \vec{u} \cdot \vec{w}$ . (distributive properties)

$\vec{u} \cdot (\vec{v} - \vec{w}) = \vec{u} \cdot \vec{v} - \vec{u} \cdot \vec{w}$ .

Let  $\vec{u}$  and  $\vec{v}$  be two vectors in  $\mathbb{R}^3$  (or  $\mathbb{R}^2$ ). There is a unique angle  $\theta$  between  $\vec{u}$  and  $\vec{v}$  with  $0 \leq \theta \leq \pi$ .



Let  $\vec{u}$  and  $\vec{v}$  be two vectors in  $\mathbb{R}^3$  (or  $\mathbb{R}^2$ ). There is a unique angle  $\theta$  between  $\vec{u}$  and  $\vec{v}$  with  $0 \leq \theta \leq \pi$ .




### Theorem

Let  $\vec{u}$  and  $\vec{v}$  be nonzero vectors, and let  $\theta$  denote the angle between  $\vec{u}$  and  $\vec{v}$ . Then

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta.$$

Let  $\vec{u}$  and  $\vec{v}$  be two vectors in  $\mathbb{R}^3$  (or  $\mathbb{R}^2$ ). There is a unique angle  $\theta$  between  $\vec{u}$  and  $\vec{v}$  with  $0 \leq \theta \leq \pi$ .



### Theorem


Let  $\vec{u}$  and  $\vec{v}$  be nonzero vectors, and let  $\theta$  denote the angle between  $\vec{u}$  and  $\vec{v}$ . Then

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta.$$



Proof.

We first prove the **Law of Cosines** – a generalization of the Pythagorean theorem:



$$\begin{aligned}c^2 &= p^2 + (b - q)^2 = a^2 \sin^2 \theta + (b - a \cos \theta)^2 \\&= a^2 (\sin^2 \theta + \cos^2 \theta) + b^2 - 2ab \cos \theta \\&= a^2 + b^2 - 2ab \cos \theta.\end{aligned}$$





### Proof. (continued)

In terms of vectors, we see that



$$\begin{aligned} \|\vec{v} - \vec{w}\|^2 &= \|\vec{v}\|^2 + \|\vec{w}\|^2 - 2\|\vec{v}\| \|\vec{w}\| \cos \theta \\ \parallel \end{aligned}$$

$$\begin{aligned} (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) &= \|\vec{v}\|^2 - 2\vec{v} \cdot \vec{w} + \|\vec{w}\|^2 \\ \Downarrow \end{aligned}$$

$$\begin{aligned} \|\vec{v}\|^2 + \|\vec{w}\|^2 - 2\|\vec{v}\| \|\vec{w}\| \cos \theta &= \|\vec{v}\|^2 - 2\vec{v} \cdot \vec{w} + \|\vec{w}\|^2 \\ \Downarrow \end{aligned}$$

## Proof. (continued)

In terms of vectors, we see that



$$\begin{aligned} \|\vec{v} - \vec{w}\|^2 &= \|\vec{v}\|^2 + \|\vec{w}\|^2 - 2\|\vec{v}\| \|\vec{w}\| \cos \theta \\ \parallel \end{aligned}$$

$$\begin{aligned} (\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) &= \|\vec{v}\|^2 - 2\vec{v} \cdot \vec{w} + \|\vec{w}\|^2 \\ \Downarrow \end{aligned}$$

$$\begin{aligned} \|\vec{v}\|^2 + \|\vec{w}\|^2 - 2\|\vec{v}\| \|\vec{w}\| \cos \theta &= \|\vec{v}\|^2 - 2\vec{v} \cdot \vec{w} + \|\vec{w}\|^2 \\ \Downarrow \end{aligned}$$

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta.$$





$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta.$$

► If  $0 \leq \theta < \frac{\pi}{2}$ , then  $\cos \theta > 0$ .

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta.$$

- If  $0 \leq \theta < \frac{\pi}{2}$ , then  $\cos \theta > 0$ .
- If  $\theta = \frac{\pi}{2}$ , then  $\cos \theta = 0$ .



$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta.$$

- ▶ If  $0 \leq \theta < \frac{\pi}{2}$ , then  $\cos \theta > 0$ .
- ▶ If  $\theta = \frac{\pi}{2}$ , then  $\cos \theta = 0$ .
- ▶ If  $\frac{\pi}{2} < \theta \leq \pi$ , then  $\cos \theta < 0$ .

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta.$$

- ▶ If  $0 \leq \theta < \frac{\pi}{2}$ , then  $\cos \theta > 0$ .
- ▶ If  $\theta = \frac{\pi}{2}$ , then  $\cos \theta = 0$ .
- ▶ If  $\frac{\pi}{2} < \theta \leq \pi$ , then  $\cos \theta < 0$ .

Therefore, for nonzero vectors  $\vec{u}$  and  $\vec{v}$ ,

- ▶  $\vec{u} \cdot \vec{v} > 0$  if and only if  $0 \leq \theta < \frac{\pi}{2}$ .

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta.$$

- ▶ If  $0 \leq \theta < \frac{\pi}{2}$ , then  $\cos \theta > 0$ .
- ▶ If  $\theta = \frac{\pi}{2}$ , then  $\cos \theta = 0$ .
- ▶ If  $\frac{\pi}{2} < \theta \leq \pi$ , then  $\cos \theta < 0$ .

Therefore, for nonzero vectors  $\vec{u}$  and  $\vec{v}$ ,

- ▶  $\vec{u} \cdot \vec{v} > 0$  if and only if  $0 \leq \theta < \frac{\pi}{2}$ .
- ▶  $\vec{u} \cdot \vec{v} = 0$  if and only if  $\theta = \frac{\pi}{2}$ .

$$\vec{u} \cdot \vec{v} = \|\vec{u}\| \|\vec{v}\| \cos \theta.$$

- ▶ If  $0 \leq \theta < \frac{\pi}{2}$ , then  $\cos \theta > 0$ .
- ▶ If  $\theta = \frac{\pi}{2}$ , then  $\cos \theta = 0$ .
- ▶ If  $\frac{\pi}{2} < \theta \leq \pi$ , then  $\cos \theta < 0$ .

Therefore, for nonzero vectors  $\vec{u}$  and  $\vec{v}$ ,

- ▶  $\vec{u} \cdot \vec{v} > 0$  if and only if  $0 \leq \theta < \frac{\pi}{2}$ .
- ▶  $\vec{u} \cdot \vec{v} = 0$  if and only if  $\theta = \frac{\pi}{2}$ .
- ▶  $\vec{u} \cdot \vec{v} < 0$  if and only if  $\frac{\pi}{2} < \theta \leq \pi$ .

## Definition

Vectors  $\vec{u}$  and  $\vec{v}$  are **orthogonal** if and only if  $\vec{u} = \vec{0}$  or  $\vec{v} = \vec{0}$  or  $\theta = \frac{\pi}{2}$ .

## Definition

Vectors  $\vec{u}$  and  $\vec{v}$  are **orthogonal** if and only if  $\vec{u} = \vec{0}$  or  $\vec{v} = \vec{0}$  or  $\theta = \frac{\pi}{2}$ .

## Theorem

Vectors  $\vec{u}$  and  $\vec{v}$  are orthogonal if and only if  $\vec{u} \cdot \vec{v} = 0$ .

## Problem

Find the angle between  $\vec{u} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$ .

## Problem

Find the angle between  $\vec{u} = \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} 0 \\ 1 \\ -1 \end{bmatrix}$ .

## Solution

$$\vec{u} \cdot \vec{v} = 1, \|\vec{u}\| = \sqrt{2} \text{ and } \|\vec{v}\| = \sqrt{2}.$$

Therefore,

$$\cos \theta = \frac{\vec{u} \cdot \vec{v}}{\|\vec{u}\| \|\vec{v}\|} = \frac{1}{\sqrt{2}\sqrt{2}} = \frac{1}{2}.$$

Since  $0 \leq \theta \leq \pi$ ,  $\theta = \frac{\pi}{3}$ .

Therefore, the angle between  $\vec{u}$  and  $\vec{v}$  is  $\frac{\pi}{3}$ .



### Problem

Find the angle between  $\vec{u} = \begin{bmatrix} 7 \\ -1 \\ 3 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} 1 \\ 4 \\ -1 \end{bmatrix}$ .

### Solution

$\vec{u} \cdot \vec{v} = 0$ , and therefore the angle between the vectors is  $\frac{\pi}{2}$ .



### Problem

Find all vectors  $\vec{v} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$  orthogonal to both  $\vec{u} = \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$  and  $\vec{w} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$ .

### Solution

There are infinitely many such vectors.

### Problem

Find all vectors  $\vec{v} = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$  orthogonal to both  $\vec{u} = \begin{bmatrix} -1 \\ -3 \\ 2 \end{bmatrix}$  and  $\vec{w} = \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$ .

### Solution

There are infinitely many such vectors. Since  $\vec{v}$  is orthogonal to both  $\vec{u}$  and  $\vec{w}$ ,

$$\begin{aligned}\vec{v} \cdot \vec{u} &= -x - 3y + 2z = 0 \\ \vec{v} \cdot \vec{w} &= y + z = 0\end{aligned}$$

### Solution (continued)

This is a homogeneous system of two linear equation in three variables.

$$\left[ \begin{array}{ccc|c} -1 & -3 & 2 & 0 \\ 0 & 1 & 1 & 0 \end{array} \right] \rightarrow \cdots \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & -5 & 0 \\ 0 & 1 & 1 & 0 \end{array} \right]$$

Therefore,  $\vec{v} = t \begin{bmatrix} 5 \\ -1 \\ 1 \end{bmatrix}$  for all  $t \in \mathbb{R}$ .

### Problem

Are  $A(4, -7, 9)$ ,  $B(6, 4, 4)$  and  $C(7, 10, -6)$  the vertices of a right angle triangle?

### Problem

Are  $A(4, -7, 9)$ ,  $B(6, 4, 4)$  and  $C(7, 10, -6)$  the vertices of a right angle triangle?

### Solution

$$\overrightarrow{AB} = \begin{bmatrix} 2 \\ 11 \\ -5 \end{bmatrix}, \quad \overrightarrow{AC} = \begin{bmatrix} 3 \\ 17 \\ -15 \end{bmatrix}, \quad \overrightarrow{BC} = \begin{bmatrix} 1 \\ 6 \\ -10 \end{bmatrix}$$

### Problem

Are  $A(4, -7, 9)$ ,  $B(6, 4, 4)$  and  $C(7, 10, -6)$  the vertices of a right angle triangle?

### Solution

$$\overrightarrow{AB} = \begin{bmatrix} 2 \\ 11 \\ -5 \end{bmatrix}, \quad \overrightarrow{AC} = \begin{bmatrix} 3 \\ 17 \\ -15 \end{bmatrix}, \quad \overrightarrow{BC} = \begin{bmatrix} 1 \\ 6 \\ -10 \end{bmatrix}$$

$$\blacktriangleright \overrightarrow{AB} \cdot \overrightarrow{AC} = 6 + 187 + 75 \neq 0.$$



## Problem

Are  $A(4, -7, 9)$ ,  $B(6, 4, 4)$  and  $C(7, 10, -6)$  the vertices of a right angle triangle?

## Solution

$$\overrightarrow{AB} = \begin{bmatrix} 2 \\ 11 \\ -5 \end{bmatrix}, \quad \overrightarrow{AC} = \begin{bmatrix} 3 \\ 17 \\ -15 \end{bmatrix}, \quad \overrightarrow{BC} = \begin{bmatrix} 1 \\ 6 \\ -10 \end{bmatrix}$$

- ▶  $\overrightarrow{AB} \cdot \overrightarrow{AC} = 6 + 187 + 75 \neq 0.$
- ▶  $\overrightarrow{BA} \cdot \overrightarrow{BC} = (-\overrightarrow{AB}) \cdot \overrightarrow{BC} = -2 - 66 - 50 \neq 0.$

## Problem

Are  $A(4, -7, 9)$ ,  $B(6, 4, 4)$  and  $C(7, 10, -6)$  the vertices of a right angle triangle?

## Solution

$$\overrightarrow{AB} = \begin{bmatrix} 2 \\ 11 \\ -5 \end{bmatrix}, \quad \overrightarrow{AC} = \begin{bmatrix} 3 \\ 17 \\ -15 \end{bmatrix}, \quad \overrightarrow{BC} = \begin{bmatrix} 1 \\ 6 \\ -10 \end{bmatrix}$$

- ▶  $\overrightarrow{AB} \cdot \overrightarrow{AC} = 6 + 187 + 75 \neq 0.$
- ▶  $\overrightarrow{BA} \cdot \overrightarrow{BC} = (-\overrightarrow{AB}) \cdot \overrightarrow{BC} = -2 - 66 - 50 \neq 0.$
- ▶  $\overrightarrow{CA} \cdot \overrightarrow{CB} = (-\overrightarrow{AC}) \cdot (-\overrightarrow{BC}) = \overrightarrow{AC} \cdot \overrightarrow{BC} = 3 + 102 + 150 \neq 0.$

### Problem

Are  $A(4, -7, 9)$ ,  $B(6, 4, 4)$  and  $C(7, 10, -6)$  the vertices of a right angle triangle?

### Solution

$$\overrightarrow{AB} = \begin{bmatrix} 2 \\ 11 \\ -5 \end{bmatrix}, \quad \overrightarrow{AC} = \begin{bmatrix} 3 \\ 17 \\ -15 \end{bmatrix}, \quad \overrightarrow{BC} = \begin{bmatrix} 1 \\ 6 \\ -10 \end{bmatrix}$$

- ▶  $\overrightarrow{AB} \cdot \overrightarrow{AC} = 6 + 187 + 75 \neq 0.$
- ▶  $\overrightarrow{BA} \cdot \overrightarrow{BC} = (-\overrightarrow{AB}) \cdot \overrightarrow{BC} = -2 - 66 - 50 \neq 0.$
- ▶  $\overrightarrow{CA} \cdot \overrightarrow{CB} = (-\overrightarrow{AC}) \cdot (-\overrightarrow{BC}) = \overrightarrow{AC} \cdot \overrightarrow{BC} = 3 + 102 + 150 \neq 0.$

Because none of the angles is  $\frac{\pi}{2}$ , the triangle is not a right angle triangle.


## Problem

A rhombus is a parallelogram with sides of equal length. Prove that the diagonals of a rhombus are perpendicular.

## Problem

A rhombus is a parallelogram with sides of equal length. Prove that the diagonals of a rhombus are perpendicular.

## Solution



Define the parallelogram (rhombus) by vectors  $\vec{u}$  and  $\vec{v}$ .


Then the diagonals are  $\vec{u} + \vec{v}$  and  $\vec{u} - \vec{v}$ .

Show that  $\vec{u} + \vec{v}$  and  $\vec{u} - \vec{v}$  are perpendicular.

## Problem

A rhombus is a parallelogram with sides of equal length. Prove that the diagonals of a rhombus are perpendicular.

## Solution



Define the parallelogram (rhombus) by vectors  $\vec{u}$  and  $\vec{v}$ .

Then the diagonals are  $\vec{u} + \vec{v}$  and  $\vec{u} - \vec{v}$ .

Show that  $\vec{u} + \vec{v}$  and  $\vec{u} - \vec{v}$  are perpendicular.

$$\begin{aligned}(\vec{u} + \vec{v}) \cdot (\vec{u} - \vec{v}) &= \vec{u} \cdot \vec{u} - \vec{u} \cdot \vec{v} + \vec{v} \cdot \vec{u} - \vec{v} \cdot \vec{v} \\&= \|\vec{u}\|^2 - \vec{u} \cdot \vec{v} + \vec{u} \cdot \vec{v} - \|\vec{v}\|^2 \\&= \|\vec{u}\|^2 - \|\vec{v}\|^2 \\&= 0, \quad \text{since } \|\vec{u}\| = \|\vec{v}\|.\end{aligned}$$

Therefore, the diagonals are perpendicular.





# Projections

Given two nonzero vectors  $\vec{u}$  and  $\vec{d}$ , one can always express  $\vec{u}$  as a sum  $\vec{u} = \vec{u}_1 + \vec{u}_2$ , where  $\vec{u}_1$  is parallel to  $\vec{d}$  and  $\vec{u}_2$  is orthogonal to  $\vec{d}$ .




# Projections

Given two nonzero vectors  $\vec{u}$  and  $\vec{d}$ , one can always express  $\vec{u}$  as a sum  $\vec{u} = \vec{u}_1 + \vec{u}_2$ , where  $\vec{u}_1$  is parallel to  $\vec{d}$  and  $\vec{u}_2$  is orthogonal to  $\vec{d}$ .



# Projections


Given two nonzero vectors  $\vec{u}$  and  $\vec{d}$ , one can always express  $\vec{u}$  as a sum  $\vec{u} = \vec{u}_1 + \vec{u}_2$ , where  $\vec{u}_1$  is parallel to  $\vec{d}$  and  $\vec{u}_2$  is orthogonal to  $\vec{d}$ .



$\vec{u}_1$  is the projection of  $\vec{u}$  onto  $\vec{d}$ , written  $\vec{u}_1 = \text{proj}_{\vec{d}} \vec{u}$ .

# Projections

Given two nonzero vectors  $\vec{u}$  and  $\vec{d}$ , one can always express  $\vec{u}$  as a sum  $\vec{u} = \vec{u}_1 + \vec{u}_2$ , where  $\vec{u}_1$  is parallel to  $\vec{d}$  and  $\vec{u}_2$  is orthogonal to  $\vec{d}$ .



$\vec{u}_1$  is the projection of  $\vec{u}$  onto  $\vec{d}$ , written  $\vec{u}_1 = \text{proj}_{\vec{d}} \vec{u}$ .

How to find  $\vec{u}_1 = \text{proj}_{\vec{d}} \vec{u}$  ?

$$\vec{u}_2 \cdot \vec{u}_1 = 0 \qquad (\vec{u}_1 \perp \vec{u}_2)$$

$$\vec{u}_2 \cdot (t\vec{d}) = 0 \qquad (\vec{u}_1 = t\vec{d})$$

$$t(\vec{u}_2 \cdot \vec{d}) = 0$$

$$\vec{u}_2 \cdot \vec{d} = 0 \qquad (t \neq 0 \text{ b.c. } \vec{u} \neq \vec{0})$$

$$(\vec{u} - \vec{u}_1) \cdot \vec{d} = 0 \qquad (\vec{u}_1 + \vec{u}_2 = \vec{u})$$

$$\vec{u} \cdot \vec{d} - \vec{u}_1 \cdot \vec{d} = 0$$

$$\vec{u} \cdot \vec{d} - (t\vec{d}) \cdot \vec{d} = 0 \qquad (\vec{u}_1 = t\vec{d})$$



$$\vec{u} \cdot \vec{d} - t(\vec{d} \cdot \vec{d}) = 0$$



$$\vec{u} \cdot \vec{d} - t||\vec{d}||^2 = 0$$

$$\vec{u} \cdot \vec{d} = t||\vec{d}||^2$$

$$t = \frac{\vec{u} \cdot \vec{d}}{||\vec{d}||^2} \qquad (\vec{d} \neq \vec{0})$$

$$\vec{u}_1 = \frac{\vec{u} \cdot \vec{d}}{||\vec{d}||^2} \vec{d} \qquad (\vec{u}_1 = t\vec{d})$$







## Theorem

Let  $\vec{u}$  and  $\vec{d}$  be vectors with  $\vec{d} \neq \vec{0}$ .

1. The projection of  $\vec{u}$  onto  $\vec{d}$  is

$$\vec{u}_1 = \text{proj}_{\vec{d}} \vec{u} = \frac{\vec{u} \cdot \vec{d}}{\|\vec{d}\|^2} \vec{d}.$$



## Theorem

Let  $\vec{u}$  and  $\vec{d}$  be vectors with  $\vec{d} \neq \vec{0}$ .

1. The projection of  $\vec{u}$  onto  $\vec{d}$  is

$$\vec{u}_1 = \text{proj}_{\vec{d}} \vec{u} = \frac{\vec{u} \cdot \vec{d}}{\|\vec{d}\|^2} \vec{d}.$$

- 2.

$$\vec{u}_2 = \vec{u} - \frac{\vec{u} \cdot \vec{d}}{\|\vec{d}\|^2} \vec{d}$$

is orthogonal to  $\vec{d}$ .

$$\vec{u}_1 = \text{proj}_{\vec{d}} \vec{u} = \left( \vec{u} \cdot \frac{\vec{d}}{||\vec{d}||} \right) \frac{\vec{d}}{||\vec{d}||} = \frac{\vec{u} \cdot \vec{d}}{||\vec{d}||^2} \vec{d}$$



unit vector



$$\vec{u}_1 = \text{proj}_{\vec{d}} \vec{u} = \left( \vec{u} \cdot \frac{\vec{d}}{||\vec{d}||} \right) \frac{\vec{d}}{||\vec{d}||} = \frac{\vec{u} \cdot \vec{d}}{||\vec{d}||^2} \vec{d}$$

unit vector



$$\vec{u}_1 = \text{proj}_{\vec{d}} \vec{u} = \left( \vec{u} \cdot \frac{\vec{d}}{\|\vec{d}\|} \right) \frac{\vec{d}}{\|\vec{d}\|} = \frac{\vec{u} \cdot \vec{d}}{\|\vec{d}\|^2} \vec{d}$$



length

unit vector



$$\vec{u}_1 = \text{proj}_{\vec{d}} \vec{u} = \left( \vec{u} \cdot \frac{\vec{d}}{||\vec{d}||} \right) \frac{\vec{d}}{||\vec{d}||} = \frac{\vec{u} \cdot \vec{d}}{||\vec{d}||^2} \vec{d}$$



length



direction

### Problem

Let  $\vec{u} = \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} 3 \\ 1 \\ -1 \end{bmatrix}$ . Find vectors  $\vec{u}_1$  and  $\vec{u}_2$  so that  $\vec{u} = \vec{u}_1 + \vec{u}_2$ , with  $\vec{u}_1$  parallel to  $\vec{v}$  and  $\vec{u}_2$  orthogonal to  $\vec{v}$ .

### Problem

Let  $\vec{u} = \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} 3 \\ 1 \\ -1 \end{bmatrix}$ . Find vectors  $\vec{u}_1$  and  $\vec{u}_2$  so that  $\vec{u} = \vec{u}_1 + \vec{u}_2$ , with  $\vec{u}_1$  parallel to  $\vec{v}$  and  $\vec{u}_2$  orthogonal to  $\vec{v}$ .

### Solution

$$\vec{u}_1 = \text{proj}_{\vec{v}} \vec{u} = \frac{\vec{u} \cdot \vec{v}}{||\vec{v}||^2} \vec{v} = \frac{5}{11} \begin{bmatrix} 3 \\ 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 15/11 \\ 5/11 \\ -5/11 \end{bmatrix}.$$

## Problem

Let  $\vec{u} = \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} 3 \\ 1 \\ -1 \end{bmatrix}$ . Find vectors  $\vec{u}_1$  and  $\vec{u}_2$  so that  $\vec{u} = \vec{u}_1 + \vec{u}_2$ , with  $\vec{u}_1$  parallel to  $\vec{v}$  and  $\vec{u}_2$  orthogonal to  $\vec{v}$ .

## Solution

$$\vec{u}_1 = \text{proj}_{\vec{v}} \vec{u} = \frac{\vec{u} \cdot \vec{v}}{||\vec{v}||^2} \vec{v} = \frac{5}{11} \begin{bmatrix} 3 \\ 1 \\ -1 \end{bmatrix} = \begin{bmatrix} 15/11 \\ 5/11 \\ -5/11 \end{bmatrix}.$$

$$\vec{u}_2 = \vec{u} - \vec{u}_1 = \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix} - \frac{5}{11} \begin{bmatrix} 3 \\ 1 \\ -1 \end{bmatrix} = \frac{1}{11} \begin{bmatrix} 7 \\ -16 \\ 5 \end{bmatrix} = \begin{bmatrix} 7/11 \\ -16/11 \\ 5/11 \end{bmatrix}.$$

## Problem

Let  $P(3, 2, -1)$  be a point in  $\mathbb{R}^3$  and  $L$  a line with equation

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} + t \begin{bmatrix} 3 \\ -1 \\ -2 \end{bmatrix}.$$

Find the **shortest distance** from  $P$  to  $L$ , and **find the point**  $Q$  on  $L$  that is closest to  $P$ .


## Problem

Let  $P(3, 2, -1)$  be a point in  $\mathbb{R}^3$  and  $L$  a line with equation

$$\begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} + t \begin{bmatrix} 3 \\ -1 \\ -2 \end{bmatrix}.$$

Find the **shortest distance** from  $P$  to  $L$ , and **find the point**  $Q$  on  $L$  that is closest to  $P$ .

## Solution



Let  $P_0 = P_0(2, 1, 3)$  be a point on  $L$ ,

and let  $\vec{d} = \begin{bmatrix} 3 & -1 & -2 \end{bmatrix}^T$ .

Then  $\overrightarrow{P_0Q} = \text{proj}_{\vec{d}} \overrightarrow{P_0P}$ ,  $\overrightarrow{OQ} = \overrightarrow{OP_0} + \overrightarrow{P_0Q}$ ,

and the shortest distance from  $P$  to  $L$  is

the length of  $\overrightarrow{QP}$ , where  $\overrightarrow{QP} = \overrightarrow{P_0P} - \overrightarrow{P_0Q}$ .



Solution (continued)

$$\overrightarrow{P_0P} = \begin{bmatrix} 1 & 1 & -4 \end{bmatrix}^T, \vec{d} = \begin{bmatrix} 3 & -1 & -2 \end{bmatrix}^T.$$

$$\overrightarrow{P_0Q} = \text{proj}_{\vec{d}} \overrightarrow{P_0P} = \frac{\overrightarrow{P_0P} \cdot \vec{d}}{\|\vec{d}\|^2} \vec{d} = \frac{10}{14} \begin{bmatrix} 3 \\ -1 \\ -2 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 15 \\ -5 \\ -10 \end{bmatrix}.$$

Solution (continued)

$$\overrightarrow{P_0P} = \begin{bmatrix} 1 & 1 & -4 \end{bmatrix}^T, \vec{d} = \begin{bmatrix} 3 & -1 & -2 \end{bmatrix}^T.$$

$$\overrightarrow{P_0Q} = \text{proj}_{\vec{d}} \overrightarrow{P_0P} = \frac{\overrightarrow{P_0P} \cdot \vec{d}}{\|\vec{d}\|^2} \vec{d} = \frac{10}{14} \begin{bmatrix} 3 \\ -1 \\ -2 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 15 \\ -5 \\ -10 \end{bmatrix}.$$

Therefore,

$$\overrightarrow{OQ} = \overrightarrow{OP_0} + \overrightarrow{P_0Q} = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix} + \frac{1}{7} \begin{bmatrix} 15 \\ -5 \\ -10 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 29 \\ 2 \\ 11 \end{bmatrix},$$

$$\text{so } Q = Q\left(\frac{29}{7}, \frac{2}{7}, \frac{11}{7}\right).$$

### Solution (continued)

Finally, the shortest distance from  $P(3, 2, -1)$  to  $L$  is the length of  $\overrightarrow{QP}$ , where

$$\overrightarrow{QP} = \overrightarrow{P_0P} - \overrightarrow{P_0Q} = \begin{bmatrix} 1 \\ 1 \\ -4 \end{bmatrix} - \frac{1}{7} \begin{bmatrix} 15 \\ -5 \\ -10 \end{bmatrix} = \frac{2}{7} \begin{bmatrix} -4 \\ 6 \\ -9 \end{bmatrix}.$$

### Solution (continued)

Finally, the shortest distance from  $P(3, 2, -1)$  to  $L$  is the length of  $\overrightarrow{QP}$ , where

$$\overrightarrow{QP} = \overrightarrow{P_0P} - \overrightarrow{P_0Q} = \begin{bmatrix} 1 \\ 1 \\ -4 \end{bmatrix} - \frac{1}{7} \begin{bmatrix} 15 \\ -5 \\ -10 \end{bmatrix} = \frac{2}{7} \begin{bmatrix} -4 \\ 6 \\ -9 \end{bmatrix}.$$

Therefore the shortest distance from  $P$  to  $L$  is


$$\|\overrightarrow{QP}\| = \frac{2}{7} \sqrt{(-4)^2 + 6^2 + (-9)^2} = \frac{2}{7} \sqrt{133}.$$



# Planes

## Definition

A nonzero vector  $\vec{n}$  is a **normal vector** to a plane if and only if  $\vec{n} \cdot \vec{v} = 0$  for every vector  $\vec{v}$  in the plane.



Given a point  $P_0$  and a nonzero vector  $\vec{n}$ , there is a unique plane containing  $P_0$  and orthogonal to  $\vec{n}$ .

Consider a plane containing a point  $P_0$  and orthogonal to vector  $\vec{n}$ , and let  $P$  be an arbitrary point on this plane.

Consider a plane containing a point  $P_0$  and orthogonal to vector  $\vec{n}$ , and let  $P$  be an arbitrary point on this plane.

Then

$$\vec{n} \cdot \overrightarrow{P_0P} = 0,$$



Consider a plane containing a point  $P_0$  and orthogonal to vector  $\vec{n}$ , and let  $P$  be an arbitrary point on this plane.

Then

$$\vec{n} \cdot \overrightarrow{P_0P} = 0,$$

or, equivalently,

$$\vec{n} \cdot (\overrightarrow{OP} - \overrightarrow{OP_0}) = 0,$$

and is a **vector equation** of the plane.









$$\vec{n} \cdot (\overrightarrow{0P} - \overrightarrow{0P_0}) = 0 \quad \Longleftrightarrow \quad \vec{n} \cdot \overrightarrow{0P} = \vec{n} \cdot \overrightarrow{0P_0}$$

by setting  $P_0 = P_0(x_0, y_0, z_0)$ ,  $P = P(x, y, z)$ ,  $\vec{n} = \begin{bmatrix} a & b & c \end{bmatrix}^T$

$$\Longleftrightarrow \quad \begin{bmatrix} a \\ b \\ c \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix} \cdot \begin{bmatrix} x_0 \\ y_0 \\ z_0 \end{bmatrix}$$

$$\Longleftrightarrow \quad ax + by + cz = ax_0 + by_0 + cz_0,$$

setting  $d = ax_0 + by_0 + cz_0$  - a scalar

$$\vec{n} \cdot (\overrightarrow{0P} - \overrightarrow{0P_0}) = 0 \quad \Longleftrightarrow \quad \vec{n} \cdot \overrightarrow{0P} = \vec{n} \cdot \overrightarrow{0P_0}$$

by setting  $P_0 = P_0(x_0, y_0, z_0)$ ,  $P = P(x, y, z)$ ,  $\vec{n} = \begin{bmatrix} a & b & c \end{bmatrix}^T$

$$\Longleftrightarrow \quad \begin{bmatrix} a \\ b \\ c \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix} \cdot \begin{bmatrix} x_0 \\ y_0 \\ z_0 \end{bmatrix}$$

$$\Longleftrightarrow \quad ax + by + cz = ax_0 + by_0 + cz_0,$$

setting  $d = ax_0 + by_0 + cz_0$  - a scalar

$$\Longleftrightarrow \quad \boxed{ax + by + cz = d}, \text{ where } a, b, c, d \in \mathbb{R}.$$

$$\vec{n} \cdot (\vec{OP} - \vec{OP_0}) = 0 \quad \Longleftrightarrow \quad \vec{n} \cdot \vec{OP} = \vec{n} \cdot \vec{OP_0}$$

by setting  $P_0 = P_0(x_0, y_0, z_0)$ ,  $P = P(x, y, z)$ ,  $\vec{n} = \begin{bmatrix} a & b & c \end{bmatrix}^T$

$$\Longleftrightarrow \begin{bmatrix} a \\ b \\ c \end{bmatrix} \cdot \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \end{bmatrix} \cdot \begin{bmatrix} x_0 \\ y_0 \\ z_0 \end{bmatrix}$$

$$\Longleftrightarrow ax + by + cz = ax_0 + by_0 + cz_0,$$

setting  $d = ax_0 + by_0 + cz_0$  - a scalar

$$\Longleftrightarrow \boxed{ax + by + cz = d}, \text{ where } a, b, c, d \in \mathbb{R}.$$

This is the **scalar equation** of the plane.



### Problem

Find an equation of the plane containing  $P_0(1, -1, 0)$  and orthogonal to  $\vec{n} = \begin{bmatrix} -3 & 5 & 2 \end{bmatrix}^T$ .

### Problem

Find an equation of the plane containing  $P_0(1, -1, 0)$  and orthogonal to  $\vec{n} = \begin{bmatrix} -3 & 5 & 2 \end{bmatrix}^T$ .

### Solution

A **vector equation** of this plane is

$$\begin{bmatrix} -3 \\ 5 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} x - 1 \\ y + 1 \\ z \end{bmatrix} = 0.$$

## Problem

Find an equation of the plane containing  $P_0(1, -1, 0)$  and orthogonal to  $\vec{n} = \begin{bmatrix} -3 & 5 & 2 \end{bmatrix}^T$ .

## Solution

A **vector equation** of this plane is

$$\begin{bmatrix} -3 \\ 5 \\ 2 \end{bmatrix} \cdot \begin{bmatrix} x - 1 \\ y + 1 \\ z \end{bmatrix} = 0.$$

A **scalar equation** of this plane is

$$-3x + 5y + 2z = -3(1) + 5(-1) + 2(0) = -8,$$

i.e., the plane has scalar equation

$$-3x + 5y + 2z = -8.$$


## Problem

Find the shortest distance from the point  $P(2, 3, 0)$  to the plane with equation  $5x + y + z = -1$ , and find the point  $Q$  on the plane that is closest to  $P$ .

## Problem

Find the shortest distance from the point  $P(2, 3, 0)$  to the plane with equation  $5x + y + z = -1$ , and find the point  $Q$  on the plane that is closest to  $P$ .

## Solution



Pick an arbitrary point  $P_0$  on the plane.

Then  $\vec{QP} = \text{proj}_{\vec{n}} \vec{P_0P}$ ,


$\|\vec{QP}\|$  is the shortest distance,

and  $\vec{OQ} = \vec{OP} - \vec{QP}$ .

## Problem

Find the shortest distance from the point  $P(2, 3, 0)$  to the plane with equation  $5x + y + z = -1$ , and find the point  $Q$  on the plane that is closest to  $P$ .

## Solution



Pick an arbitrary point  $P_0$  on the plane.

Then  $\overrightarrow{QP} = \text{proj}_{\vec{n}} \overrightarrow{P_0P}$ ,

$\|\overrightarrow{QP}\|$  is the shortest distance,


and  $\overrightarrow{OQ} = \overrightarrow{OP} - \overrightarrow{QP}$ .

$$\vec{n} = \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T.$$

## Problem

Find the shortest distance from the point  $P(2, 3, 0)$  to the plane with equation  $5x + y + z = -1$ , and find the point  $Q$  on the plane that is closest to  $P$ .

## Solution



Pick an arbitrary point  $P_0$  on the plane.

Then  $\overrightarrow{QP} = \text{proj}_{\vec{n}} \overrightarrow{P_0P}$ ,

$\|\overrightarrow{QP}\|$  is the shortest distance,


and  $\overrightarrow{OQ} = \overrightarrow{OP} - \overrightarrow{QP}$ .

$\vec{n} = \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T$ . Choose  $P_0 = P_0(0, 0, -1)$ .

## Problem

Find the shortest distance from the point  $P(2, 3, 0)$  to the plane with equation  $5x + y + z = -1$ , and find the point  $Q$  on the plane that is closest to  $P$ .

## Solution



Pick an arbitrary point  $P_0$  on the plane.

Then  $\overrightarrow{QP} = \text{proj}_{\vec{n}} \overrightarrow{P_0P}$ ,

$\|\overrightarrow{QP}\|$  is the shortest distance,


and  $\overrightarrow{OQ} = \overrightarrow{OP} - \overrightarrow{QP}$ .

$\vec{n} = \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T$ . Choose  $P_0 = P_0(0, 0, -1)$ . Then

$$\overrightarrow{P_0P} = \begin{bmatrix} 2 & 3 & 1 \end{bmatrix}^T$$




Solution (continued)



$$\overrightarrow{P_0P} = \begin{bmatrix} 2 & 3 & 1 \end{bmatrix}^T.$$

$$\vec{n} = \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T.$$

### Solution (continued)




$$\overrightarrow{P_0P} = \begin{bmatrix} 2 & 3 & 1 \end{bmatrix}^T.$$

$$\vec{n} = \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T.$$

$$\overrightarrow{QP} = \text{proj}_{\vec{n}} \overrightarrow{P_0P} = \frac{\overrightarrow{P_0P} \cdot \vec{n}}{\|\vec{n}\|^2} \vec{n} = \frac{14}{27} \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T.$$

# Solution (continued)




$$\overrightarrow{P_0P} = \begin{bmatrix} 2 & 3 & 1 \end{bmatrix}^T.$$

$$\vec{n} = \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T.$$

$$\overrightarrow{QP} = \text{proj}_{\vec{n}} \overrightarrow{P_0P} = \frac{\overrightarrow{P_0P} \cdot \vec{n}}{\|\vec{n}\|^2} \vec{n} = \frac{14}{27} \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T.$$

Since  $\|\overrightarrow{QP}\| = \frac{14}{27} \sqrt{27} = \frac{14\sqrt{3}}{9}$ , the shortest distance from  $P$  to the plane is  $\frac{14\sqrt{3}}{9}$ .

# Solution (continued)



$$\overrightarrow{P_0P} = \begin{bmatrix} 2 & 3 & 1 \end{bmatrix}^T.$$

$$\vec{n} = \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T.$$


$$\overrightarrow{QP} = \text{proj}_{\vec{n}} \overrightarrow{P_0P} = \frac{\overrightarrow{P_0P} \cdot \vec{n}}{\|\vec{n}\|^2} \vec{n} = \frac{14}{27} \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T.$$

Since  $\|\overrightarrow{QP}\| = \frac{14}{27}\sqrt{27} = \frac{14\sqrt{3}}{9}$ , the shortest distance from P to the plane is  $\frac{14\sqrt{3}}{9}$ .

To find Q, we have

$$\begin{aligned} \overrightarrow{OQ} = \overrightarrow{OP} - \overrightarrow{QP} &= \begin{bmatrix} 2 & 3 & 0 \end{bmatrix}^T - \frac{14}{27} \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T \\ &= \frac{1}{27} \begin{bmatrix} -16 & 67 & -14 \end{bmatrix}^T. \end{aligned}$$

## Solution (continued)



$$\overrightarrow{P_0P} = \begin{bmatrix} 2 & 3 & 1 \end{bmatrix}^T.$$

$$\vec{n} = \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T.$$

$$\overrightarrow{QP} = \text{proj}_{\vec{n}} \overrightarrow{P_0P} = \frac{\overrightarrow{P_0P} \cdot \vec{n}}{\|\vec{n}\|^2} \vec{n} = \frac{14}{27} \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T.$$

Since  $\|\overrightarrow{QP}\| = \frac{14}{27}\sqrt{27} = \frac{14\sqrt{3}}{9}$ , the shortest distance from P to the plane is  $\frac{14\sqrt{3}}{9}$ .

To find Q, we have

$$\begin{aligned} \overrightarrow{OQ} = \overrightarrow{OP} - \overrightarrow{QP} &= \begin{bmatrix} 2 & 3 & 0 \end{bmatrix}^T - \frac{14}{27} \begin{bmatrix} 5 & 1 & 1 \end{bmatrix}^T \\ &= \frac{1}{27} \begin{bmatrix} -16 & 67 & -14 \end{bmatrix}^T. \end{aligned}$$

Therefore  $Q = Q \left( -\frac{16}{27}, \frac{67}{27}, -\frac{14}{27} \right)$ .

### Remark

Here is a general answer: the distance from  $P(x_0, y_0, z_0)$  to the plane  $ax + by + cz = d$  is

$$\text{distance} = \frac{|ax_0 + by_0 + cz_0 - d|}{\sqrt{a^2 + b^2 + c^2}}$$



# The Cross Product

## Definition

Let  $\vec{u} = \begin{bmatrix} x_1 & y_1 & z_1 \end{bmatrix}^T$  and  $\vec{v} = \begin{bmatrix} x_2 & y_2 & z_2 \end{bmatrix}^T$ . Then

$$\vec{u} \times \vec{v} = \begin{bmatrix} y_1 z_2 - z_1 y_2 \\ -(x_1 z_2 - z_1 x_2) \\ x_1 y_2 - y_1 x_2 \end{bmatrix}.$$




# The Cross Product

## Definition

Let  $\vec{u} = \begin{bmatrix} x_1 & y_1 & z_1 \end{bmatrix}^T$  and  $\vec{v} = \begin{bmatrix} x_2 & y_2 & z_2 \end{bmatrix}^T$ . Then

$$\vec{u} \times \vec{v} = \begin{bmatrix} y_1 z_2 - z_1 y_2 \\ -(x_1 z_2 - z_1 x_2) \\ x_1 y_2 - y_1 x_2 \end{bmatrix}.$$



## Remark

$\vec{u} \times \vec{v}$  is a vector:

- Direction: orthogonal to both  $\vec{u}$  and  $\vec{v}$ .
- Size: the area of the corresponding parallelogram.







## Theorem

Let  $\vec{v}, \vec{w} \in \mathbb{R}^3$ .

1.  $\vec{v} \times \vec{w}$  is orthogonal to both  $\vec{v}$  and  $\vec{w}$ .
2. If  $\vec{v}$  and  $\vec{w}$  are both nonzero, then  $\vec{v} \times \vec{w} = \vec{0}$  if and only if  $\vec{v}$  and  $\vec{w}$  are parallel.

## Problem

Find all vectors orthogonal to both  $\vec{u} = \begin{bmatrix} -1 & -3 & 2 \end{bmatrix}^T$  and  $\vec{v} = \begin{bmatrix} 0 & 1 & 1 \end{bmatrix}^T$ . (We previously solved this using the **dot product**.)

## Problem

Find all vectors orthogonal to both  $\vec{u} = \begin{bmatrix} -1 & -3 & 2 \end{bmatrix}^T$  and  $\vec{v} = \begin{bmatrix} 0 & 1 & 1 \end{bmatrix}^T$ . (We previously solved this using the **dot product**.)

## Solution

$$\vec{u} \times \vec{v} = \begin{vmatrix} \vec{i} & -1 & 0 \\ \vec{j} & -3 & 1 \\ \vec{k} & 2 & 1 \end{vmatrix} = -5\vec{i} + \vec{j} - \vec{k} = \begin{bmatrix} -5 \\ 1 \\ -1 \end{bmatrix}.$$

Any scalar multiple of  $\vec{u} \times \vec{v}$  is also orthogonal to both  $\vec{u}$  and  $\vec{v}$ , so

$$t \begin{bmatrix} -5 \\ 1 \\ -1 \end{bmatrix}, \quad \forall t \in \mathbb{R},$$

gives all vectors orthogonal to both  $\vec{u}$  and  $\vec{v}$ .

(Compare this with our earlier answer.)

## Problem


Given two lines

$$L_1 : \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ -1 \end{bmatrix} + s \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \quad \text{and} \quad L_2 : \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} + t \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix},$$

- A. Find the shortest distance between  $L_1$  and  $L_2$ .
- B. Find the points P on  $L_1$  and Q on  $L_2$  that are closest together.




## Solution



Choose  $P_1(3, 1, -1)$  on  $L_1$  and  $P_2(1, 2, 0)$  on  $L_2$ .


Let  $\vec{d}_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}$  and  $\vec{d}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$  denote direction

vectors for  $L_1$  and  $L_2$ , respectively.



$$\vec{d}_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \vec{d}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$$


The shortest distance between  $L_1$  and  $L_2$  is the length of the projection of  $\overrightarrow{P_1P_2}$  onto  $\vec{n} = \vec{d}_1 \times \vec{d}_2$ .



$$\vec{d}_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \vec{d}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$$

The shortest distance between  $L_1$  and  $L_2$  is the length of the projection of  $\overrightarrow{P_1P_2}$  onto  $\vec{n} = \vec{d}_1 \times \vec{d}_2$ .

$$\overrightarrow{P_1P_2} = \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix} \quad \text{and} \quad \vec{n} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \times \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix}$$




$$\vec{d}_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \vec{d}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$$

The shortest distance between  $L_1$  and  $L_2$  is the length of the projection of  $\overrightarrow{P_1P_2}$  onto  $\vec{n} = \vec{d}_1 \times \vec{d}_2$ .

$$\overrightarrow{P_1P_2} = \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix} \quad \text{and} \quad \vec{n} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \times \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix}$$

$$\text{proj}_{\vec{n}} \overrightarrow{P_1P_2} = \frac{\overrightarrow{P_1P_2} \cdot \vec{n}}{\|\vec{n}\|^2} \vec{n}, \quad \text{and} \quad \|\text{proj}_{\vec{n}} \overrightarrow{P_1P_2}\| = \frac{|\overrightarrow{P_1P_2} \cdot \vec{n}|}{\|\vec{n}\|}.$$



$$\vec{d}_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \vec{d}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$$


The shortest distance between  $L_1$  and  $L_2$  is the length of the projection of  $\overrightarrow{P_1P_2}$  onto  $\vec{n} = \vec{d}_1 \times \vec{d}_2$ .

$$\overrightarrow{P_1P_2} = \begin{bmatrix} -2 \\ 1 \\ 1 \end{bmatrix} \quad \text{and} \quad \vec{n} = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix} \times \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} = \begin{bmatrix} 2 \\ -3 \\ -1 \end{bmatrix}$$

$$\text{proj}_{\vec{n}} \overrightarrow{P_1P_2} = \frac{\overrightarrow{P_1P_2} \cdot \vec{n}}{\|\vec{n}\|^2} \vec{n}, \quad \text{and} \quad \|\text{proj}_{\vec{n}} \overrightarrow{P_1P_2}\| = \frac{|\overrightarrow{P_1P_2} \cdot \vec{n}|}{\|\vec{n}\|}.$$

Therefore, the shortest distance between  $L_1$  and  $L_2$  is  $\frac{|-8|}{\sqrt{14}} = \frac{4}{7}\sqrt{14}$ .

Solution B.




$$\vec{d}_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \vec{d}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix};$$

$$\vec{OP} = \begin{bmatrix} 3 + s \\ 1 + s \\ -1 - s \end{bmatrix} \text{ for some } s \in \mathbb{R};$$

$$\vec{OQ} = \begin{bmatrix} 1 + t \\ 2 \\ 2t \end{bmatrix} \text{ for some } t \in \mathbb{R}.$$

Solution B.



$$\vec{d}_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \vec{d}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix};$$


$$\vec{OP} = \begin{bmatrix} 3+s \\ 1+s \\ -1-s \end{bmatrix} \text{ for some } s \in \mathbb{R};$$

$$\vec{OQ} = \begin{bmatrix} 1+t \\ 2 \\ 2t \end{bmatrix} \text{ for some } t \in \mathbb{R}.$$

Now  $\vec{PQ} = \begin{bmatrix} -2-s+t & 1-s & 1+s+2t \end{bmatrix}^T$  is orthogonal to both  $L_1$  and  $L_2$ , so

$$\vec{PQ} \cdot \vec{d}_1 = 0 \quad \text{and} \quad \vec{PQ} \cdot \vec{d}_2 = 0,$$

Solution B.



$$\vec{d}_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \vec{d}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix};$$

$$\vec{OP} = \begin{bmatrix} 3+s \\ 1+s \\ -1-s \end{bmatrix} \text{ for some } s \in \mathbb{R};$$

$$\vec{OQ} = \begin{bmatrix} 1+t \\ 2 \\ 2t \end{bmatrix} \text{ for some } t \in \mathbb{R}.$$

Now  $\vec{PQ} = \begin{bmatrix} -2-s+t & 1-s & 1+s+2t \end{bmatrix}^T$  is orthogonal to both  $L_1$  and  $L_2$ , so


$$\vec{PQ} \cdot \vec{d}_1 = 0 \quad \text{and} \quad \vec{PQ} \cdot \vec{d}_2 = 0,$$

i.e.,

$$\begin{aligned} -2 - 3s - t &= 0 \\ s + 5t &= 0. \end{aligned}$$



Solution B.



$$\vec{d}_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \vec{d}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix};$$

$$\vec{OP} = \begin{bmatrix} 3+s \\ 1+s \\ -1-s \end{bmatrix} \text{ for some } s \in \mathbb{R};$$

$$\vec{OQ} = \begin{bmatrix} 1+t \\ 2 \\ 2t \end{bmatrix} \text{ for some } t \in \mathbb{R}.$$

Now  $\vec{PQ} = \begin{bmatrix} -2-s+t & 1-s & 1+s+2t \end{bmatrix}^T$  is orthogonal to both  $L_1$  and  $L_2$ , so


$$\vec{PQ} \cdot \vec{d}_1 = 0 \quad \text{and} \quad \vec{PQ} \cdot \vec{d}_2 = 0,$$

i.e.,

$$\begin{aligned} -2 - 3s - t &= 0 \\ s + 5t &= 0. \end{aligned}$$

This system has unique solution  $s = -\frac{5}{7}$  and  $t = \frac{1}{7}$ .

Solution B.



$$\vec{d}_1 = \begin{bmatrix} 1 \\ 1 \\ -1 \end{bmatrix}, \vec{d}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix};$$

$$\vec{OP} = \begin{bmatrix} 3+s \\ 1+s \\ -1-s \end{bmatrix} \text{ for some } s \in \mathbb{R};$$

$$\vec{OQ} = \begin{bmatrix} 1+t \\ 2 \\ 2t \end{bmatrix} \text{ for some } t \in \mathbb{R}.$$

Now  $\vec{PQ} = \begin{bmatrix} -2-s+t & 1-s & 1+s+2t \end{bmatrix}^T$  is orthogonal to both  $L_1$  and  $L_2$ , so

$$\vec{PQ} \cdot \vec{d}_1 = 0 \quad \text{and} \quad \vec{PQ} \cdot \vec{d}_2 = 0,$$

i.e.,

$$\begin{aligned} -2-3s-t &= 0 \\ s+5t &= 0. \end{aligned}$$

This system has unique solution  $s = -\frac{5}{7}$  and  $t = \frac{1}{7}$ . Therefore,

$$P = P\left(\frac{16}{7}, \frac{2}{7}, -\frac{2}{7}\right) \quad \text{and} \quad Q = Q\left(\frac{8}{7}, 2, \frac{2}{7}\right).$$

The shortest distance between  $L_1$  and  $L_2$  is  $\|\overrightarrow{PQ}\|$ . Since

$$P = P\left(\frac{16}{7}, \frac{2}{7}, -\frac{2}{7}\right) \quad \text{and} \quad Q = Q\left(\frac{8}{7}, 2, \frac{2}{7}\right),$$

The shortest distance between  $L_1$  and  $L_2$  is  $||\overrightarrow{PQ}||$ . Since

$$P = P\left(\frac{16}{7}, \frac{2}{7}, -\frac{2}{7}\right) \quad \text{and} \quad Q = Q\left(\frac{8}{7}, 2, \frac{2}{7}\right),$$

$$\overrightarrow{PQ} = \frac{1}{7} \begin{bmatrix} 8 \\ 14 \\ 2 \end{bmatrix} - \frac{1}{7} \begin{bmatrix} 16 \\ 2 \\ -2 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} -8 \\ 12 \\ 4 \end{bmatrix},$$

The shortest distance between  $L_1$  and  $L_2$  is  $\|\overrightarrow{PQ}\|$ . Since

$$P = P\left(\frac{16}{7}, \frac{2}{7}, -\frac{2}{7}\right) \quad \text{and} \quad Q = Q\left(\frac{8}{7}, 2, \frac{2}{7}\right),$$

$$\overrightarrow{PQ} = \frac{1}{7} \begin{bmatrix} 8 \\ 14 \\ 2 \end{bmatrix} - \frac{1}{7} \begin{bmatrix} 16 \\ 2 \\ -2 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} -8 \\ 12 \\ 4 \end{bmatrix},$$

and

$$\|\overrightarrow{PQ}\| = \frac{1}{7}\sqrt{224} = \frac{4}{7}\sqrt{14}.$$

The shortest distance between  $L_1$  and  $L_2$  is  $||\overrightarrow{PQ}||$ . Since

$$P = P\left(\frac{16}{7}, \frac{2}{7}, -\frac{2}{7}\right) \quad \text{and} \quad Q = Q\left(\frac{8}{7}, 2, \frac{2}{7}\right),$$

$$\overrightarrow{PQ} = \frac{1}{7} \begin{bmatrix} 8 \\ 14 \\ 2 \end{bmatrix} - \frac{1}{7} \begin{bmatrix} 16 \\ 2 \\ -2 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} -8 \\ 12 \\ 4 \end{bmatrix},$$

and

$$||\overrightarrow{PQ}|| = \frac{1}{7}\sqrt{224} = \frac{4}{7}\sqrt{14}.$$

Therefore the shortest distance between  $L_1$  and  $L_2$  is  $\frac{4}{7}\sqrt{14}$ .



## Shortest Distances

### Problem ( Challenge Problem )

Write yourself a plan to find the shortest distance in  $\mathbb{R}^3$  between either a point, line or plane, to either a point, line or plane.





## Point-point distance

If  $P$  and  $Q$  are two points, then  $d(P, Q) = |\overrightarrow{PQ}|$ .






## Point-plane distance

If  $P$  is a point and  $\Sigma : \vec{n} \cdot \vec{x} = d$  is a plane containing a point  $Q$ , then

$$d(P, \Sigma) = \frac{|\overrightarrow{PQ} \cdot \vec{n}|}{|\vec{n}|}$$






## Point-line distance

If  $P$  is a point and  $L$  is a line  $\vec{r}(t) = Q + t\vec{u}$ , then

$$d(P, L) = \frac{|\vec{PQ} \times \vec{u}|}{|\vec{u}|}$$






## Line-line distance

If  $L$  is a line  $\vec{r}(t) = \vec{Q} + t\vec{u}$  and  $M$  is another line  $\vec{s} = \vec{P} + t\vec{v}$ , then

$$d(L, M) = \frac{|\overrightarrow{PQ} \cdot (\vec{u} \times \vec{v})|}{|\vec{u} \times \vec{v}|}$$







## Plane-plane distance

If  $\Sigma : \vec{n} \cdot \vec{x} = d$  and  $\Theta : \vec{n} \cdot \vec{x} = e$  are two parallel planes, then

$$d(\Sigma, \Theta) = \frac{|e - d|}{|\vec{n}|}$$

