

# Math 221: LINEAR ALGEBRA

## Chapter 6. Vector Spaces §6-3. Linear Independence

Le Chen<sup>1</sup>

Emory University, 2020 Fall

(last updated on 11/03/2020)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.

Linear Independence

The Fundamental Theorem

Bases and Dimension



# Linear Independence

## Definition

Let  $V$  be a vector space and  $S = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$  a subset of  $V$ . The set  $S$  is **linearly independent** or simply **independent** if the following condition holds:

$$s_1\mathbf{u}_1 + s_2\mathbf{u}_2 + \cdots + s_k\mathbf{u}_k = \mathbf{0} \quad \Rightarrow \quad s_1 = s_2 = \cdots = s_k = 0$$

# Linear Independence

## Definition

Let  $V$  be a vector space and  $S = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$  a subset of  $V$ . The set  $S$  is **linearly independent** or simply **independent** if the following condition holds:

$$s_1\mathbf{u}_1 + s_2\mathbf{u}_2 + \cdots + s_k\mathbf{u}_k = \mathbf{0} \quad \Rightarrow \quad s_1 = s_2 = \cdots = s_k = 0$$

i.e., the only linear combination that vanishes is the trivial one.

# Linear Independence

## Definition

Let  $V$  be a vector space and  $S = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$  a subset of  $V$ . The set  $S$  is **linearly independent** or simply **independent** if the following condition holds:

$$s_1\mathbf{u}_1 + s_2\mathbf{u}_2 + \cdots + s_k\mathbf{u}_k = \mathbf{0} \quad \Rightarrow \quad s_1 = s_2 = \cdots = s_k = 0$$

i.e., the only linear combination that vanishes is the trivial one. If  $S$  is not linearly independent, then  $S$  is said to be **dependent**.

### Example

The set  $S = \left\{ \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} \right\}$  is a **dependent** subset of  $\mathbb{R}^3$

because

$$a \begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix} + b \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} + c \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

has nontrivial solutions, for example  $a = 2$ ,  $b = 3$  and  $c = -1$ .

## Problem

Is the set  $T = \{3x^2 - x + 2, x^2 + x - 1, x^2 - 3x + 4\}$  an independent subset of  $\mathcal{P}_2$ ?



## Problem

Is the set  $T = \{3x^2 - x + 2, x^2 + x - 1, x^2 - 3x + 4\}$  an independent subset of  $\mathcal{P}_2$ ?

## Solution

Suppose  $a(3x^2 - x + 2) + b(x^2 + x - 1) + c(x^2 - 3x + 4) = 0$ , for some  $a, b, c \in \mathbb{R}$ . Then

$$x^2(3a + b + c) + x(-a + b - 3c) + (2a - b + 4c) = 0,$$

implying that

$$\begin{aligned} 3a + b + c &= 0 \\ -a + b - 3c &= 0 \\ 2a - b + 4c &= 0 \end{aligned}$$

## Problem

Is the set  $T = \{3x^2 - x + 2, x^2 + x - 1, x^2 - 3x + 4\}$  an independent subset of  $\mathcal{P}_2$ ?

## Solution

Suppose  $a(3x^2 - x + 2) + b(x^2 + x - 1) + c(x^2 - 3x + 4) = 0$ , for some  $a, b, c \in \mathbb{R}$ . Then

$$x^2(3a + b + c) + x(-a + b - 3c) + (2a - b + 4c) = 0,$$

implying that

$$\begin{aligned} 3a + b + c &= 0 \\ -a + b - 3c &= 0 \\ 2a - b + 4c &= 0 \end{aligned}$$

Solving this linear system of three equations in three variables

$$\left[ \begin{array}{ccc|c} 3 & 1 & 1 & 0 \\ -1 & 1 & -3 & 0 \\ 2 & -1 & 4 & 0 \end{array} \right] \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & 1 & 0 \\ 0 & 1 & -2 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right].$$

Since there is nontrivial solution,  $T$  is a dependent subset of  $\mathcal{P}_2$ . ■

## Problem

Is  $U = \left\{ \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \right\}$  an independent subset of  $\mathbf{M}_{22}$ ?

## Problem

Is  $U = \left\{ \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \right\}$  an independent subset of  $\mathbf{M}_{22}$ ?

## Solution

Suppose  $a \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + c \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$  for some  $a, b, c \in \mathbb{R}$ .

## Problem

Is  $U = \left\{ \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \right\}$  an independent subset of  $\mathbf{M}_{22}$ ?

## Solution

Suppose  $a \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + c \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$  for some

$a, b, c \in \mathbb{R}$ .

↓

$$\begin{array}{rcl} a + c & = & 0, \quad a + b = 0, \\ b + c & = & 0, \quad a + c = 0. \end{array}$$

## Problem

Is  $U = \left\{ \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \right\}$  an independent subset of  $\mathbf{M}_{22}$ ?

## Solution

Suppose  $a \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + c \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$  for some  $a, b, c \in \mathbb{R}$ .

↓

$$\begin{array}{rcl} a + c & = & 0, \quad a + b = 0, \\ b + c & = & 0, \quad a + c = 0. \end{array}$$

This system of four equations in three variables has **unique solution**  
 $a = b = c = 0$ ,

## Problem

Is  $U = \left\{ \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \right\}$  an independent subset of  $\mathbf{M}_{22}$ ?

## Solution

Suppose  $a \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} + b \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + c \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$  for some  $a, b, c \in \mathbb{R}$ .

↓

$$\begin{array}{rcl} a + c & = & 0, & a + b & = & 0, \\ b + c & = & 0, & a + c & = & 0. \end{array}$$

This system of four equations in three variables has **unique solution**  $a = b = c = 0$ ,

↓

$U$  is an independent subset of  $\mathbf{M}_{22}$ .



### Example (An independent subset of $\mathcal{P}_n$ )

Consider  $\{1, x, x^2, \dots, x^n\}$ , and suppose that

$$a_0 \cdot 1 + a_1x + a_2x^2 + \cdots + a_nx^n = 0$$

for some  $a_0, a_1, \dots, a_n \in \mathbb{R}$ . Then  $a_0 = a_1 = \cdots = a_n = 0$ , and thus  $\{1, x, x^2, \dots, x^n\}$  is an independent subset of  $\mathcal{P}_n$ .



### Example (An independent subset of $\mathcal{P}_n$ )

Consider  $\{1, x, x^2, \dots, x^n\}$ , and suppose that

$$a_0 \cdot 1 + a_1x + a_2x^2 + \cdots + a_nx^n = 0$$

for some  $a_0, a_1, \dots, a_n \in \mathbb{R}$ . Then  $a_0 = a_1 = \cdots = a_n = 0$ , and thus  $\{1, x, x^2, \dots, x^n\}$  is an independent subset of  $\mathcal{P}_n$ .

### Example ( Polynomials with distinct degrees )

Any set of polynomials with DISTINCT degrees is independent.

For example,

$$\{2x^4 - x^3 + 5, \quad -3x^3 + 2x^2 + 2, \quad 4x^2 + x - 3, \quad 2x - 1, \quad 3\}$$

is an independent subset of  $\mathcal{P}_4$ .

## Example

As we saw earlier,  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  (the standard basis of  $\mathbb{R}^n$ ) is an independent subset of  $\mathbb{R}^n$ .

## Example

As we saw earlier,  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  (the standard basis of  $\mathbb{R}^n$ ) is an independent subset of  $\mathbb{R}^n$ .

## Example

$$U = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$$

is an independent subset of  $\mathbf{M}_{32}$ .

## Example

As we saw earlier,  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  (the standard basis of  $\mathbb{R}^n$ ) is an independent subset of  $\mathbb{R}^n$ .

## Example

$$U = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}$$

is an independent subset of  $\mathbf{M}_{32}$ .

## Example ( An independent subset of $\mathbf{M}_{mn}$ )

In general, the set of  $mn$   $m \times n$  matrices that have a '1' in position  $(i, j)$  and zeros elsewhere,  $1 \leq i \leq m$ ,  $1 \leq j \leq n$ , constitutes an independent subset of  $\mathbf{M}_{mn}$ .

## Example

Let  $V$  be a vector space.

1. If  $\mathbf{v}$  is a **nonzero** vector of  $V$ , then  $\{\mathbf{v}\}$  is an independent subset of  $V$ .

## Example

Let  $V$  be a vector space.

1. If  $\mathbf{v}$  is a **nonzero** vector of  $V$ , then  $\{\mathbf{v}\}$  is an independent subset of  $V$ .

**Proof.** Suppose that  $k\mathbf{v} = \mathbf{0}$  for some  $k \in \mathbb{R}$ . Since  $\mathbf{v} \neq \mathbf{0}$ , it must be that  $k = 0$ , and therefore  $\{\mathbf{v}\}$  is an independent set. ■

2. The zero vector of  $V$ ,  $\mathbf{0}$  is never an element of an independent subset of  $V$ .

## Example

Let  $V$  be a vector space.

1. If  $\mathbf{v}$  is a **nonzero** vector of  $V$ , then  $\{\mathbf{v}\}$  is an independent subset of  $V$ .

**Proof.** Suppose that  $k\mathbf{v} = \mathbf{0}$  for some  $k \in \mathbb{R}$ . Since  $\mathbf{v} \neq \mathbf{0}$ , it must be that  $k = 0$ , and therefore  $\{\mathbf{v}\}$  is an independent set. ■

2. The zero vector of  $V$ ,  $\mathbf{0}$  is never an element of an independent subset of  $V$ .

**Proof.** Suppose  $S = \{\mathbf{0}, \mathbf{v}_2, \mathbf{v}_3, \dots, \mathbf{v}_k\}$  is a subset of  $V$ . Then

$$1(\mathbf{0}) + 0(\mathbf{v}_2) + 0(\mathbf{v}_3) + \cdots + 0(\mathbf{v}_k) = \mathbf{0}.$$

Since the coefficient of  $\mathbf{0}$  (on the left-hand side) is '1', we have a nontrivial vanishing linear combination of the vectors of  $S$ . Therefore,  $S$  is dependent. ■

## Problem

Let  $V$  be a vector space and let  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  be an independent subset of  $V$ . Is

$$S = \{\mathbf{u} + \mathbf{v}, 2\mathbf{u} + \mathbf{w}, \mathbf{v} - 5\mathbf{w}\}$$

an independent subset of  $V$ ? Justify your answer.



## Problem

Let  $V$  be a vector space and let  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  be an independent subset of  $V$ . Is

$$S = \{\mathbf{u} + \mathbf{v}, 2\mathbf{u} + \mathbf{w}, \mathbf{v} - 5\mathbf{w}\}$$

an independent subset of  $V$ ? Justify your answer.

## Solution

Suppose that a linear combination of the vectors of  $S$  is equal to zero, i.e.,

$$a(\mathbf{u} + \mathbf{v}) + b(2\mathbf{u} + \mathbf{w}) + c(\mathbf{v} - 5\mathbf{w}) = \mathbf{0}$$

for some  $a, b, c \in \mathbb{R}$ . Then  $(a + 2b)\mathbf{u} + (a + c)\mathbf{v} + (b - 5c)\mathbf{w} = \mathbf{0}$ . Since  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  is independent,

$$a + 2b = 0$$

$$a + c = 0$$

$$b - 5c = 0.$$

Solving for  $a, b$  and  $c$ , we find that the system has unique solution  $a = b = c = 0$ . Therefore,  $S$  is linearly independent. ■

## Problem

Suppose that  $A$  is an  $n \times n$  matrix with the property that  $A^k = \mathbf{0}$  but  $A^{k-1} \neq \mathbf{0}$ . Prove that

$$B = \{I, A, A^2, \dots, A^{k-1}\}$$

is an independent subset of  $\mathbf{M}_{nn}$ .

## Problem

Suppose that  $A$  is an  $n \times n$  matrix with the property that  $A^k = \mathbf{0}$  but  $A^{k-1} \neq \mathbf{0}$ . Prove that

$$B = \{I, A, A^2, \dots, A^{k-1}\}$$

is an independent subset of  $\mathbf{M}_{nn}$ .

## Solution ( hints only )

Use the standard approach: take a linear combination of the matrices and set it equal to the  $n \times n$  zero matrix. Two key points:

- ▶ Since  $A^{k-1} \neq \mathbf{0}$ , the matrices  $A, A^2, \dots, A^{k-2}$  are all nonzero.
- ▶ Since  $A^k = \mathbf{0}$ , the matrices  $A^{k+1}, A^{k+2}, A^{k+3}, \dots$  are all zero.

### Theorem (Unique Representation Theorem)

Let  $V$  be a vector space and let  $U = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\} \subseteq V$  be an independent set. If  $\mathbf{v}$  is in  $\text{span}(U)$ , then  $\mathbf{v}$  has a unique representation as a linear combination of elements of  $U$ .

### Theorem (Unique Representation Theorem)

Let  $V$  be a vector space and let  $U = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\} \subseteq V$  be an independent set. If  $\mathbf{v}$  is in  $\text{span}(U)$ , then  $\mathbf{v}$  has a unique representation as a linear combination of elements of  $U$ .

Again, the proof of the corresponding result for  $\mathbb{R}^n$  generalizes to an arbitrary vector space  $V$ .



## The Fundamental Theorem

The Fundamental Theorem for  $\mathbb{R}^n$  generalizes to an arbitrary vector space.

## The Fundamental Theorem

The Fundamental Theorem for  $\mathbb{R}^n$  generalizes to an arbitrary vector space.

### Theorem (Fundamental Theorem)

Let  $V$  be a vector space that can be spanned by a set of  $n$  vectors, and suppose that  $V$  contains an independent subset of  $m$  vectors. Then  $m \leq n$ .





Proof. (continued)

Let  $A = [ a_{ij} ]$ , and suppose that  $m > n$ . Since  $\text{rank}(A) = \dim(\text{row}(A)) \leq n$ , it follows that the rows of  $A$  form a dependent subset of  $\mathbb{R}^n$ , and hence there is a nontrivial linear combination of the rows of  $A$  that is equal to the  $1 \times n$  vector of all zeros, i.e., there exist  $s_1, s_2, \dots, s_m \in \mathbb{R}$ , not all equal to zero, such that

$$[ s_1 \quad s_2 \quad \cdots \quad s_m ] A = [ 0 \quad 0 \quad \cdots \quad 0 ] = \mathbf{0}_{1n}.$$

It follows that for each  $j$ ,  $1 \leq j \leq n$ ,

$$s_1 a_{1j} + s_2 a_{2j} + \cdots + s_m a_{mj} = 0. \tag{1}$$

Consider the (nontrivial) linear combination of vectors of  $Y$ :

$$s_1 \mathbf{y}_1 + s_2 \mathbf{y}_2 + \cdots + s_m \mathbf{y}_m.$$

Proof. (continued)

$$\begin{aligned} s_1 \mathbf{y}_1 + s_2 \mathbf{y}_2 + \cdots + s_m \mathbf{y}_m &= s_1 (a_{11} \mathbf{x}_1 + a_{12} \mathbf{x}_2 + \cdots + a_{1n} \mathbf{x}_n) + \\ &\quad s_2 (a_{21} \mathbf{x}_1 + a_{22} \mathbf{x}_2 + \cdots + a_{2n} \mathbf{x}_n) + \\ &\quad \vdots \\ &\quad s_m (a_{m1} \mathbf{x}_1 + a_{m2} \mathbf{x}_2 + \cdots + a_{mn} \mathbf{x}_n) \\ &= (s_1 a_{11} + s_2 a_{21} + \cdots + s_m a_{m1}) \mathbf{x}_1 + \\ &\quad (s_1 a_{12} + s_2 a_{22} + \cdots + s_m a_{m2}) \mathbf{x}_2 + \\ &\quad \vdots \\ &\quad (s_1 a_{1n} + s_2 a_{2n} + \cdots + s_m a_{mn}) \mathbf{x}_n. \end{aligned}$$

By Equation (1), it follows that

$$s_1 \mathbf{y}_1 + s_2 \mathbf{y}_2 + \cdots + s_m \mathbf{y}_m = 0 \mathbf{x}_1 + 0 \mathbf{x}_2 + \cdots + 0 \mathbf{x}_n = \mathbf{0}.$$

Therefore,  $s_1 \mathbf{y}_1 + s_2 \mathbf{y}_2 + \cdots + s_m \mathbf{y}_m = \mathbf{0}$  is a nontrivial vanishing linear combination of the vectors of  $Y$ .

Proof. (continued)

$$\begin{aligned} s_1 \mathbf{y}_1 + s_2 \mathbf{y}_2 + \cdots + s_m \mathbf{y}_m &= s_1 (a_{11} \mathbf{x}_1 + a_{12} \mathbf{x}_2 + \cdots + a_{1n} \mathbf{x}_n) + \\ &\quad s_2 (a_{21} \mathbf{x}_1 + a_{22} \mathbf{x}_2 + \cdots + a_{2n} \mathbf{x}_n) + \\ &\quad \vdots \\ &\quad s_m (a_{m1} \mathbf{x}_1 + a_{m2} \mathbf{x}_2 + \cdots + a_{mn} \mathbf{x}_n) \\ &= (s_1 a_{11} + s_2 a_{21} + \cdots + s_m a_{m1}) \mathbf{x}_1 + \\ &\quad (s_1 a_{12} + s_2 a_{22} + \cdots + s_m a_{m2}) \mathbf{x}_2 + \\ &\quad \vdots \\ &\quad (s_1 a_{1n} + s_2 a_{2n} + \cdots + s_m a_{mn}) \mathbf{x}_n. \end{aligned}$$

By Equation (1), it follows that

$$s_1 \mathbf{y}_1 + s_2 \mathbf{y}_2 + \cdots + s_m \mathbf{y}_m = 0 \mathbf{x}_1 + 0 \mathbf{x}_2 + \cdots + 0 \mathbf{x}_n = \mathbf{0}.$$

Therefore,  $s_1 \mathbf{y}_1 + s_2 \mathbf{y}_2 + \cdots + s_m \mathbf{y}_m = \mathbf{0}$  is a nontrivial vanishing linear combination of the vectors of  $Y$ . This contradicts the fact that  $Y$  is independent, and therefore  $m \leq n$ . ■



# Bases and Dimension

## Definition

Let  $V$  be a vector space and let  $B = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\} \subseteq V$ . We say  $B$  is a **basis** of  $V$  if

- (i)  $B$  is an independent subset of  $V$  and
- (ii)  $\text{span}(B) = V$ .

# Bases and Dimension

## Definition

Let  $V$  be a vector space and let  $B = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\} \subseteq V$ . We say  $B$  is a **basis** of  $V$  if

- (i)  $B$  is an independent subset of  $V$  and
- (ii)  $\text{span}(B) = V$ .

## Remark (Unique Representation Theorem)

Recall that if  $V$  is a vector space and  $B$  is a basis of  $V$ , then as seen earlier, any vector  $\mathbf{u} \in V$  can be expressed uniquely as a linear combination of vectors of  $B$ .

## Example

As we saw earlier,  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  is a basis of  $\mathbb{R}^n$ , called the standard basis of  $\mathbb{R}^n$ .



## Example

As we saw earlier,  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  is a basis of  $\mathbb{R}^n$ , called the standard basis of  $\mathbb{R}^n$ .

## Example (A basis of $\mathcal{P}_n$ )

We've already seen that

$$\{1, x, x^2, \dots, x^n\}$$

spans  $\mathcal{P}_n$  and is an independent subset of  $\mathcal{P}_n$ , and is thus a basis of  $\mathcal{P}_n$ .

## Example

As we saw earlier,  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  is a basis of  $\mathbb{R}^n$ , called the standard basis of  $\mathbb{R}^n$ .

## Example (A basis of $\mathcal{P}_n$ )

We've already seen that

$$\{1, x, x^2, \dots, x^n\}$$

spans  $\mathcal{P}_n$  and is an independent subset of  $\mathcal{P}_n$ , and is thus a basis of  $\mathcal{P}_n$ .

$$\{1, x, x^2, \dots, x^n\}$$

is called the standard basis of  $\mathcal{P}_n$ .

## Example

As we saw earlier,  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  is a basis of  $\mathbb{R}^n$ , called the standard basis of  $\mathbb{R}^n$ .

## Example (A basis of $\mathcal{P}_n$ )

We've already seen that

$$\{1, x, x^2, \dots, x^n\}$$

spans  $\mathcal{P}_n$  and is an independent subset of  $\mathcal{P}_n$ , and is thus a basis of  $\mathcal{P}_n$ .

$$\{1, x, x^2, \dots, x^n\}$$

is called the standard basis of  $\mathcal{P}_n$ .

## Example (A basis of $\mathbf{M}_{mn}$ )

The set of  $mn$   $m \times n$  matrices that have a '1' in position  $(i, j)$  and zeros elsewhere,  $1 \leq i \leq m$ ,  $1 \leq j \leq n$ , spans  $\mathbf{M}_{mn}$  and is an independent subset of  $\mathbf{M}_{mn}$ .

## Example

As we saw earlier,  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  is a basis of  $\mathbb{R}^n$ , called the standard basis of  $\mathbb{R}^n$ .

## Example (A basis of $\mathcal{P}_n$ )

We've already seen that

$$\{1, x, x^2, \dots, x^n\}$$

spans  $\mathcal{P}_n$  and is an independent subset of  $\mathcal{P}_n$ , and is thus a basis of  $\mathcal{P}_n$ .

$$\{1, x, x^2, \dots, x^n\}$$

is called the standard basis of  $\mathcal{P}_n$ .

## Example (A basis of $\mathbf{M}_{mn}$ )

The set of  $mn$   $m \times n$  matrices that have a '1' in position  $(i, j)$  and zeros elsewhere,  $1 \leq i \leq m$ ,  $1 \leq j \leq n$ , spans  $\mathbf{M}_{mn}$  and is an independent subset of  $\mathbf{M}_{mn}$ . Therefore, this set constitutes a basis of  $\mathbf{M}_{mn}$  and is called the standard basis of  $\mathbf{M}_{mn}$ .

The Invariance Theorem generalizes from  $\mathbb{R}^n$  to an arbitrary vector space  $V$ . The proof is identical, and involves two applications of the Fundamental Theorem.

The Invariance Theorem generalizes from  $\mathbb{R}^n$  to an arbitrary vector space  $V$ . The proof is identical, and involves two applications of the Fundamental Theorem.

### Theorem (Invariance Theorem)

If  $V$  is a vector space with bases  $\{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_m\}$  and  $\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_n\}$ , then  $m = n$ .

The Invariance Theorem generalizes from  $\mathbb{R}^n$  to an arbitrary vector space  $V$ . The proof is identical, and involves two applications of the Fundamental Theorem.

### Theorem (Invariance Theorem)

If  $V$  is a vector space with bases  $\{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_m\}$  and  $\{\mathbf{f}_1, \mathbf{f}_2, \dots, \mathbf{f}_n\}$ , then  $m = n$ .

### Definition (Dimension of a vector space)

Let  $V$  be a vector space and suppose  $B = \{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$  is a basis of  $V$ . The **dimension** of  $V$  is the number of vectors in  $B$ , and we write  $\dim(V) = n$ . By convention,  $\dim(\{\mathbf{0}\}) := 0$ .

## Example

Let  $V$  be a vector space and  $\mathbf{u}$  a NONZERO vector of  $V$ . Then  $U = \text{span}\{\mathbf{u}\}$  is spanned by  $\{\mathbf{u}\}$ . Since  $\{\mathbf{u}\}$  is independent,  $\{\mathbf{u}\}$  is a basis of  $U$ , and thus  $\dim(U) = 1$ .



### Example

Let  $V$  be a vector space and  $\mathbf{u}$  a NONZERO vector of  $V$ . Then  $U = \text{span}\{\mathbf{u}\}$  is spanned by  $\{\mathbf{u}\}$ . Since  $\{\mathbf{u}\}$  is independent,  $\{\mathbf{u}\}$  is a basis of  $U$ , and thus  $\dim(U) = 1$ .

### Example

Since  $\{1, x, x^2, \dots, x^n\}$  is a basis of  $\mathcal{P}_n$ ,  $\dim(\mathcal{P}_n) = n + 1$ .

### Example

Let  $V$  be a vector space and  $\mathbf{u}$  a NONZERO vector of  $V$ . Then  $U = \text{span}\{\mathbf{u}\}$  is spanned by  $\{\mathbf{u}\}$ . Since  $\{\mathbf{u}\}$  is independent,  $\{\mathbf{u}\}$  is a basis of  $U$ , and thus  $\dim(U) = 1$ .

### Example

Since  $\{1, x, x^2, \dots, x^n\}$  is a basis of  $\mathcal{P}_n$ ,  $\dim(\mathcal{P}_n) = n + 1$ .

### Example

$\dim(\mathbf{M}_{mn}) = mn$  since the standard basis of  $\mathbf{M}_{mn}$  consists of  $mn$  matrices.

## Problem

Let  $U = \left\{ A \in \mathbf{M}_{22} \mid A \begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} A \right\}$ . Then  $U$  is a subspace of  $\mathbf{M}_{22}$ . Find a basis of  $U$ , and hence  $\dim(U)$ .

## Problem

Let  $U = \left\{ A \in \mathbf{M}_{22} \mid A \begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} A \right\}$ . Then  $U$  is a subspace of  $\mathbf{M}_{22}$ . Find a basis of  $U$ , and hence  $\dim(U)$ .

## Solution

Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbf{M}_{22}$ . Then

$$A \begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} a+b & -b \\ c+d & -d \end{bmatrix}$$

and

$$\begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} A = \begin{bmatrix} 1 & 1 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a+c & b+d \\ -c & -d \end{bmatrix}.$$

If  $A \in U$ , then  $\begin{bmatrix} a+b & -b \\ c+d & -d \end{bmatrix} = \begin{bmatrix} a+c & b+d \\ -c & -d \end{bmatrix}$ .

## Solution (continued)

Equating entries leads to a system of four equations in the four variables  $a$ ,  $b$ ,  $c$  and  $d$ .

$$\begin{array}{rcl} a + b & = & a + c \\ -b & = & b + d \\ c + d & = & -c \\ -d & = & -d \end{array} \quad \text{or} \quad \begin{array}{rcl} b - c & = & 0 \\ -2b - d & = & 0 \\ 2c + d & = & 0 \end{array} .$$

## Solution (continued)

Equating entries leads to a system of four equations in the four variables  $a, b, c$  and  $d$ .

$$\begin{array}{rcl} a + b & = & a + c \\ -b & = & b + d \\ c + d & = & -c \\ -d & = & -d \end{array} \quad \text{or} \quad \begin{array}{rcl} b - c & = & 0 \\ -2b - d & = & 0 \\ 2c + d & = & 0 \end{array} .$$

The solution to this system is  $a = s$ ,  $b = -\frac{1}{2}t$ ,  $c = -\frac{1}{2}t$ ,  $d = t$  for any  $s, t \in \mathbb{R}$ , and thus  $A = \begin{bmatrix} s & \frac{t}{2} \\ -\frac{t}{2} & t \end{bmatrix}$ ,  $s, t \in \mathbb{R}$ .

## Solution (continued)

Equating entries leads to a system of four equations in the four variables  $a, b, c$  and  $d$ .

$$\begin{array}{rcl} a + b & = & a + c \\ -b & = & b + d \\ c + d & = & -c \\ -d & = & -d \end{array} \quad \text{or} \quad \begin{array}{rcl} b - c & = & 0 \\ -2b - d & = & 0 \\ 2c + d & = & 0 \end{array} .$$

The solution to this system is  $a = s$ ,  $b = -\frac{1}{2}t$ ,  $c = -\frac{1}{2}t$ ,  $d = t$  for any  $s, t \in \mathbb{R}$ , and thus  $A = \begin{bmatrix} s & \frac{t}{2} \\ -\frac{t}{2} & t \end{bmatrix}$ ,  $s, t \in \mathbb{R}$ . Since  $A \in U$  is arbitrary,

$$\begin{aligned} U &= \left\{ \begin{bmatrix} s & \frac{t}{2} \\ -\frac{t}{2} & t \end{bmatrix} \mid s, t \in \mathbb{R} \right\} \\ &= \left\{ s \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + t \begin{bmatrix} 0 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{bmatrix} \mid s, t \in \mathbb{R} \right\} \\ &= \text{span} \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{bmatrix} \right\}. \end{aligned}$$

### Solution (continued)

Let

$$B = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & -\frac{1}{2} \\ -\frac{1}{2} & 1 \end{bmatrix} \right\}.$$

Then  $\text{span}(B) = U$ , and it is routine to verify that  $B$  is an independent subset of  $\mathbf{M}_{22}$ . Therefore  $B$  is a basis of  $\mathbf{M}_{22}$ , and  $\dim(U) = 2$ . ■



## Problem

Let  $U = \{p(x) \in \mathcal{P}_2 \mid p(1) = 0\}$ . Then  $U$  is a subspace of  $\mathcal{P}_2$ . Find a basis of  $U$ , and hence  $\dim(U)$ .

## Problem

Let  $U = \{p(x) \in \mathcal{P}_2 \mid p(1) = 0\}$ . Then  $U$  is a subspace of  $\mathcal{P}_2$ . Find a basis of  $U$ , and hence  $\dim(U)$ .

## Solution

Final Answer  $B = \{x - x^2, 1 - x^2\}$  is a basis of  $U$  and thus  $\dim(U) = 2$ . ■