

# Math 221: LINEAR ALGEBRA

## Chapter 7. Linear Transformations

### §7-1. Examples and Elementary Properties

Le Chen<sup>1</sup>

Emory University, 2021 Spring

(last updated on 04/05/2021)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.

What is a Linear Transformations

Examples and Problems

Properties of Linear Transformations

Constructing Linear Transformations

## What is a Linear Transformations


Examples and Problems

Properties of Linear Transformations

Constructing Linear Transformations

What is a Linear Transformation?

## What is a Linear Transformation?



### Definition

Let  $V$  and  $W$  be vector spaces, and  $T : V \rightarrow W$  a function. Then  $T$  is called a **linear transformation** if it satisfies the following two properties.


1.  $T$  preserves addition.

For all  $\vec{v}_1, \vec{v}_2 \in V$ ,  $T(\vec{v}_1 + \vec{v}_2) = T(\vec{v}_1) + T(\vec{v}_2)$ .

2.  $T$  preserves scalar multiplication.

For all  $\vec{v} \in V$  and  $r \in \mathbb{R}$ ,  $T(r\vec{v}) = rT(\vec{v})$ .

# What is a Linear Transformation?



## Definition

Let  $V$  and  $W$  be vector spaces, and  $T : V \rightarrow W$  a function. Then  $T$  is called a **linear transformation** if it satisfies the following two properties.

1.  $T$  preserves addition.

For all  $\vec{v}_1, \vec{v}_2 \in V$ ,  $T(\vec{v}_1 + \vec{v}_2) = T(\vec{v}_1) + T(\vec{v}_2)$ .

2.  $T$  preserves scalar multiplication.

For all  $\vec{v} \in V$  and  $r \in \mathbb{R}$ ,  $T(r\vec{v}) = rT(\vec{v})$ .

## Remark

Note that the sum  $\vec{v}_1 + \vec{v}_2$  is in  $V$ , while the sum  $T(\vec{v}_1) + T(\vec{v}_2)$  is in  $W$ . Similarly,  $r\vec{v}$  is scalar multiplication in  $V$ , while  $rT(\vec{v})$  is scalar multiplication in  $W$ .

### Theorem ( Linear Transformations from $\mathbb{R}^n$ to $\mathbb{R}^m$ )

If  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is a linear transformation, then  $T$  is induced by an  $m \times n$  matrix

$$A = [ T(\vec{e}_1) \quad T(\vec{e}_2) \quad \cdots \quad T(\vec{e}_n) ],$$

where  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  is the standard basis of  $\mathbb{R}^n$ , and thus for each  $\vec{x} \in \mathbb{R}^n$

$$T(\vec{x}) = A\vec{x}.$$

## Example

$T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$  is defined by  $T \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} x + y \\ x - z \end{bmatrix}$  for all  $\begin{bmatrix} x \\ y \\ z \end{bmatrix} \in \mathbb{R}^3$ .

One can show that  $T$  preserves addition and scalar multiplication, and hence is a linear transformation. Therefore, the matrix that induces  $T$  is

$$A = \left[ T \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad T \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right] = \begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & -1 \end{bmatrix}.$$



### Remark ( Notation and Terminology )

1. If  $A$  is an  $m \times n$  matrix, then  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$  defined by

$$T_A(\vec{x}) = A\vec{x} \text{ for all } \vec{x} \in \mathbb{R}^n$$

is the linear (or matrix) transformation induced by  $A$ .

2. Let  $V$  be a vector space. A linear transformation  $T : V \rightarrow V$  is called a **linear operator on  $V$** .

What is a Linear Transformations

**Examples and Problems**

Properties of Linear Transformations

Constructing Linear Transformations

# Examples and Problems

## Example

Let  $V$  and  $W$  be vector spaces.

1. The zero transformation.

$0 : V \rightarrow W$  is defined by  $0(\vec{x}) = \vec{0}$  for all  $\vec{x} \in V$ .

# Examples and Problems

## Example

Let  $V$  and  $W$  be vector spaces.

1. The zero transformation.

$0 : V \rightarrow W$  is defined by  $0(\vec{x}) = \vec{0}$  for all  $\vec{x} \in V$ .

2. The identity operator on  $V$ .

$1_V : V \rightarrow V$  is defined by  $1_V(\vec{x}) = \vec{x}$  for all  $\vec{x} \in V$ .

# Examples and Problems

## Example

Let  $V$  and  $W$  be vector spaces.

1. The zero transformation.

$0 : V \rightarrow W$  is defined by  $0(\vec{x}) = \vec{0}$  for all  $\vec{x} \in V$ .

2. The identity operator on  $V$ .

$1_V : V \rightarrow V$  is defined by  $1_V(\vec{x}) = \vec{x}$  for all  $\vec{x} \in V$ .

3. The scalar operator on  $V$ .

Let  $a \in \mathbb{R}$ .  $s_a : V \rightarrow V$  is defined by  $s_a(\vec{x}) = a\vec{x}$  for all  $\vec{x} \in V$ .

## Problem

For vector spaces  $V$  and  $W$ , prove that the zero transformation  $0$ , the identity operator  $1_V$ , and the scalar operator  $s_a$  are linear transformations.

## Problem

For vector spaces  $V$  and  $W$ , prove that the zero transformation  $0$ , the identity operator  $1_V$ , and the scalar operator  $s_a$  are linear transformations.

## Solution ( the scalar operator )

Let  $V$  be a vector space and let  $a \in \mathbb{R}$ .

## Problem

For vector spaces  $V$  and  $W$ , prove that the zero transformation  $0$ , the identity operator  $1_V$ , and the scalar operator  $s_a$  are linear transformations.

## Solution ( the scalar operator )

Let  $V$  be a vector space and let  $a \in \mathbb{R}$ .

1. Let  $\vec{u}, \vec{w} \in V$ . Then  $s_a(\vec{u}) = a\vec{u}$  and  $s_a(\vec{w}) = a\vec{w}$ . Now

$$s_a(\vec{u} + \vec{w}) = a(\vec{u} + \vec{w}) = a\vec{u} + a\vec{w} = s_a(\vec{u}) + s_a(\vec{w}),$$

and thus  $s_a$  preserves addition.



## Problem

For vector spaces  $V$  and  $W$ , prove that the zero transformation  $0$ , the identity operator  $1_V$ , and the scalar operator  $s_a$  are linear transformations.

## Solution ( the scalar operator )

Let  $V$  be a vector space and let  $a \in \mathbb{R}$ .

1. Let  $\vec{u}, \vec{w} \in V$ . Then  $s_a(\vec{u}) = a\vec{u}$  and  $s_a(\vec{w}) = a\vec{w}$ . Now

$$s_a(\vec{u} + \vec{w}) = a(\vec{u} + \vec{w}) = a\vec{u} + a\vec{w} = s_a(\vec{u}) + s_a(\vec{w}),$$

and thus  $s_a$  preserves addition.

2. Let  $\vec{u} \in V$  and  $k \in \mathbb{R}$ . Then  $s_a(\vec{u}) = a\vec{u}$ . Now

$$s_a(k\vec{u}) = ak\vec{u} = ka\vec{u} = ks_a(\vec{u}),$$

and thus  $s_a$  preserves scalar multiplication.

## Problem

For vector spaces  $V$  and  $W$ , prove that the zero transformation  $0$ , the identity operator  $1_V$ , and the scalar operator  $s_a$  are linear transformations.

## Solution ( the scalar operator )

Let  $V$  be a vector space and let  $a \in \mathbb{R}$ .

1. Let  $\vec{u}, \vec{w} \in V$ . Then  $s_a(\vec{u}) = a\vec{u}$  and  $s_a(\vec{w}) = a\vec{w}$ . Now

$$s_a(\vec{u} + \vec{w}) = a(\vec{u} + \vec{w}) = a\vec{u} + a\vec{w} = s_a(\vec{u}) + s_a(\vec{w}),$$

and thus  $s_a$  preserves addition.

2. Let  $\vec{u} \in V$  and  $k \in \mathbb{R}$ . Then  $s_a(\vec{u}) = a\vec{u}$ . Now

$$s_a(k\vec{u}) = ak\vec{u} = ka\vec{u} = ks_a(\vec{u}),$$

and thus  $s_a$  preserves scalar multiplication.

Since  $s_a$  preserves addition and scalar multiplication,  $s_a$  is a linear transformation. ■

### Problem (Matrix transposition)

Let  $R : \mathbf{M}_{nn} \rightarrow \mathbf{M}_{nn}$  be a transformation defined by

$$R(A) = A^T \text{ for all } A \in \mathbf{M}_{nn}.$$

Show that  $R$  is a linear transformation.

### Problem (Matrix transposition)

Let  $R : \mathbf{M}_{nn} \rightarrow \mathbf{M}_{nn}$  be a transformation defined by

$$R(A) = A^T \text{ for all } A \in \mathbf{M}_{nn}.$$

Show that  $R$  is a linear transformation.

### Solution

1. Let  $A, B \in \mathbf{M}_{nn}$ . Then  $R(A) = A^T$  and  $R(B) = B^T$ , so

$$R(A + B) = (A + B)^T = A^T + B^T = R(A) + R(B).$$

## Problem (Matrix transposition)

Let  $R : \mathbf{M}_{nn} \rightarrow \mathbf{M}_{nn}$  be a transformation defined by

$$R(A) = A^T \text{ for all } A \in \mathbf{M}_{nn}.$$

Show that  $R$  is a linear transformation.

## Solution

1. Let  $A, B \in \mathbf{M}_{nn}$ . Then  $R(A) = A^T$  and  $R(B) = B^T$ , so

$$R(A + B) = (A + B)^T = A^T + B^T = R(A) + R(B).$$

2. Let  $A \in \mathbf{M}_{nn}$  and let  $k \in \mathbb{R}$ . Then  $R(A) = A^T$ , and

$$R(kA) = (kA)^T = kA^T = kR(A).$$

## Problem (Matrix transposition)

Let  $R : \mathbf{M}_{nn} \rightarrow \mathbf{M}_{nn}$  be a transformation defined by

$$R(A) = A^T \text{ for all } A \in \mathbf{M}_{nn}.$$

Show that  $R$  is a linear transformation.

## Solution

1. Let  $A, B \in \mathbf{M}_{nn}$ . Then  $R(A) = A^T$  and  $R(B) = B^T$ , so

$$R(A + B) = (A + B)^T = A^T + B^T = R(A) + R(B).$$

2. Let  $A \in \mathbf{M}_{nn}$  and let  $k \in \mathbb{R}$ . Then  $R(A) = A^T$ , and

$$R(kA) = (kA)^T = kA^T = kR(A).$$

Since  $R$  preserves addition and scalar multiplication,  $R$  is a linear transformation. ■

### Problem (Evaluation at a point)

For each  $a \in \mathbb{R}$ , the transformation  $E_a : \mathcal{P}_n \rightarrow \mathbb{R}$  is defined by

$$E_a(p) = p(a) \text{ for all } p \in \mathcal{P}_n.$$

Show that  $E_a$  is a linear transformation.

### Problem (Evaluation at a point)

For each  $a \in \mathbb{R}$ , the transformation  $E_a : \mathcal{P}_n \rightarrow \mathbb{R}$  is defined by

$$E_a(p) = p(a) \text{ for all } p \in \mathcal{P}_n.$$

Show that  $E_a$  is a linear transformation.

### Solution

1. Let  $p, q \in \mathcal{P}_n$ . Then  $E_a(p) = p(a)$  and  $E_a(q) = q(a)$ , so

$$E_a(p + q) = (p + q)(a) = p(a) + q(a) = E_a(p) + E_a(q).$$



### Problem (Evaluation at a point)

For each  $a \in \mathbb{R}$ , the transformation  $E_a : \mathcal{P}_n \rightarrow \mathbb{R}$  is defined by

$$E_a(p) = p(a) \text{ for all } p \in \mathcal{P}_n.$$

Show that  $E_a$  is a linear transformation.

### Solution

1. Let  $p, q \in \mathcal{P}_n$ . Then  $E_a(p) = p(a)$  and  $E_a(q) = q(a)$ , so

$$E_a(p + q) = (p + q)(a) = p(a) + q(a) = E_a(p) + E_a(q).$$

2. Let  $p \in \mathcal{P}_n$  and  $k \in \mathbb{R}$ . Then  $E_a(p) = p(a)$  and

$$E_a(kp) = (kp)(a) = kp(a) = kE_a(p).$$

### Problem (Evaluation at a point)

For each  $a \in \mathbb{R}$ , the transformation  $E_a : \mathcal{P}_n \rightarrow \mathbb{R}$  is defined by

$$E_a(p) = p(a) \text{ for all } p \in \mathcal{P}_n.$$

Show that  $E_a$  is a linear transformation.

### Solution

1. Let  $p, q \in \mathcal{P}_n$ . Then  $E_a(p) = p(a)$  and  $E_a(q) = q(a)$ , so

$$E_a(p + q) = (p + q)(a) = p(a) + q(a) = E_a(p) + E_a(q).$$

2. Let  $p \in \mathcal{P}_n$  and  $k \in \mathbb{R}$ . Then  $E_a(p) = p(a)$  and

$$E_a(kp) = (kp)(a) = kp(a) = kE_a(p).$$

Since  $E_a$  preserves addition and scalar multiplication,  $E_a$  is a linear transformation. ■

### Problem

Let  $S : \mathbf{M}_{nn} \rightarrow \mathbb{R}$  be a transformation defined by

$$S(A) = \text{tr}(A) \text{ for all } A \in \mathbf{M}_{nn}.$$

Prove that  $S$  is a linear transformation.

## Solution

Let  $A = [a_{ij}]$  and  $B = [b_{ij}]$  be  $n \times n$  matrices. Then

$$S(A) = \sum_{i=1}^n a_{ii} \quad \text{and} \quad S(B) = \sum_{i=1}^n b_{ii}.$$

## Solution

Let  $A = [a_{ij}]$  and  $B = [b_{ij}]$  be  $n \times n$  matrices. Then

$$S(A) = \sum_{i=1}^n a_{ii} \quad \text{and} \quad S(B) = \sum_{i=1}^n b_{ii}.$$

1. Since  $A + B = [a_{ij} + b_{ij}]$ ,

$$S(A+B) = \text{tr}(A+B) = \sum_{i=1}^n (a_{ii} + b_{ii}) = \left( \sum_{i=1}^n a_{ii} \right) + \left( \sum_{i=1}^n b_{ii} \right) = S(A) + S(B).$$

## Solution

Let  $A = [a_{ij}]$  and  $B = [b_{ij}]$  be  $n \times n$  matrices. Then

$$S(A) = \sum_{i=1}^n a_{ii} \quad \text{and} \quad S(B) = \sum_{i=1}^n b_{ii}.$$

1. Since  $A + B = [a_{ij} + b_{ij}]$ ,

$$S(A+B) = \text{tr}(A+B) = \sum_{i=1}^n (a_{ii} + b_{ii}) = \left( \sum_{i=1}^n a_{ii} \right) + \left( \sum_{i=1}^n b_{ii} \right) = S(A) + S(B).$$

2. Let  $k \in \mathbb{R}$ . Since  $kA = [ka_{ij}]$ ,

$$S(kA) = \text{tr}(kA) = \sum_{i=1}^n ka_{ii} = k \sum_{i=1}^n a_{ii} = kS(A).$$

## Solution

Let  $A = [a_{ij}]$  and  $B = [b_{ij}]$  be  $n \times n$  matrices. Then

$$S(A) = \sum_{i=1}^n a_{ii} \quad \text{and} \quad S(B) = \sum_{i=1}^n b_{ii}.$$

1. Since  $A + B = [a_{ij} + b_{ij}]$ ,

$$S(A+B) = \text{tr}(A+B) = \sum_{i=1}^n (a_{ii} + b_{ii}) = \left( \sum_{i=1}^n a_{ii} \right) + \left( \sum_{i=1}^n b_{ii} \right) = S(A) + S(B).$$

2. Let  $k \in \mathbb{R}$ . Since  $kA = [ka_{ij}]$ ,

$$S(kA) = \text{tr}(kA) = \sum_{i=1}^n ka_{ii} = k \sum_{i=1}^n a_{ii} = kS(A).$$

Therefore,  $S$  preserves addition and scalar multiplication, and thus is a linear transformation. ■

## Problem

Show that the differentiation and integration operations on  $\mathbf{P}_n$  are linear transformations. More precisely,

$$D : \mathbf{P}_n \rightarrow \mathbf{P}_{n-1} \quad \text{where } D[p(x)] = p'(x) \text{ for all } p(x) \text{ in } \mathbf{P}_n$$

$$I : \mathbf{P}_n \rightarrow \mathbf{P}_{n+1} \quad \text{where } I[p(x)] = \int_0^x p(t)dt \text{ for all } p(x) \text{ in } \mathbf{P}_n$$

are linear transformations.



## Problem

Show that the differentiation and integration operations on  $\mathbf{P}_n$  are linear transformations. More precisely,

$$D : \mathbf{P}_n \rightarrow \mathbf{P}_{n-1} \quad \text{where } D[p(x)] = p'(x) \text{ for all } p(x) \text{ in } \mathbf{P}_n$$

$$I : \mathbf{P}_n \rightarrow \mathbf{P}_{n+1} \quad \text{where } I[p(x)] = \int_0^x p(t)dt \text{ for all } p(x) \text{ in } \mathbf{P}_n$$

are linear transformations.

## Solution (Sketch)

$$[p(x) + q(x)]' = p'(x) + q'(x), \quad [rp(x)]' = (rp)'(x)$$

$$\int_0^x [p(t) + q(t)] dt = \int_0^x p(t)dt + \int_0^x q(t)dt, \quad \int_0^x rp(t)dt = r \int_0^x p(t)dt$$



What is a Linear Transformations

Examples and Problems

**Properties of Linear Transformations**

Constructing Linear Transformations

# Properties of Linear Transformations

# Properties of Linear Transformations

## Theorem

Let  $V$  and  $W$  be vector spaces, and  $T : V \rightarrow W$  a linear transformation.  
Then

1.  $T$  preserves the zero vector.  $T(\vec{0}) = \vec{0}$ .
2.  $T$  preserves additive inverses. For all  $\vec{v} \in V$ ,  $T(-\vec{v}) = -T(\vec{v})$ .
3.  $T$  preserves linear combinations.

For all  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_m \in V$  and all  $k_1, k_2, \dots, k_m \in \mathbb{R}$ ,

$$T(k_1\vec{v}_1 + k_2\vec{v}_2 + \cdots + k_m\vec{v}_m) = k_1T(\vec{v}_1) + k_2T(\vec{v}_2) + \cdots + k_mT(\vec{v}_m).$$

Proof.

1. Let  $\vec{0}_V$  denote the zero vector of  $V$  and let  $\vec{0}_W$  denote the zero vector of  $W$ . We want to prove that  $T(\vec{0}_V) = \vec{0}_W$ . Let  $\vec{x} \in V$ . Then  $0\vec{x} = \vec{0}_V$  and

$$T(\vec{0}_V) = T(0\vec{x}) = 0T(\vec{x}) = \vec{0}_W.$$

Proof.

1. Let  $\vec{0}_V$  denote the zero vector of  $V$  and let  $\vec{0}_W$  denote the zero vector of  $W$ . We want to prove that  $T(\vec{0}_V) = \vec{0}_W$ . Let  $\vec{x} \in V$ . Then  $0\vec{x} = \vec{0}_V$  and

$$T(\vec{0}_V) = T(0\vec{x}) = 0T(\vec{x}) = \vec{0}_W.$$

2. Let  $\vec{v} \in V$ ; then  $-\vec{v} \in V$  is the additive inverse of  $\vec{v}$ , so  $\vec{v} + (-\vec{v}) = \vec{0}_V$ .  
Thus

$$\begin{aligned}T(\vec{v} + (-\vec{v})) &= T(\vec{0}_V) \\T(\vec{v}) + T(-\vec{v}) &= \vec{0}_W \\T(-\vec{v}) &= \vec{0}_W - T(\vec{v}) = -T(\vec{v}).\end{aligned}$$

Proof.

1. Let  $\vec{0}_V$  denote the zero vector of  $V$  and let  $\vec{0}_W$  denote the zero vector of  $W$ . We want to prove that  $T(\vec{0}_V) = \vec{0}_W$ . Let  $\vec{x} \in V$ . Then  $0\vec{x} = \vec{0}_V$  and

$$T(\vec{0}_V) = T(0\vec{x}) = 0T(\vec{x}) = \vec{0}_W.$$

2. Let  $\vec{v} \in V$ ; then  $-\vec{v} \in V$  is the additive inverse of  $\vec{v}$ , so  $\vec{v} + (-\vec{v}) = \vec{0}_V$ .  
Thus

$$\begin{aligned}T(\vec{v} + (-\vec{v})) &= T(\vec{0}_V) \\T(\vec{v}) + T(-\vec{v}) &= \vec{0}_W \\T(-\vec{v}) &= \vec{0}_W - T(\vec{v}) = -T(\vec{v}).\end{aligned}$$

3. This result follows from preservation of addition and preservation of scalar multiplication. A formal proof would be by induction on  $m$ .



## Problem

Let  $T : \mathcal{P}_2 \rightarrow \mathbb{R}$  be a linear transformation such that

$$T(x^2 + x) = -1; \quad T(x^2 - x) = 1; \quad T(x^2 + 1) = 3.$$

Find  $T(4x^2 + 5x - 3)$ .



## Problem

Let  $T : \mathcal{P}_2 \rightarrow \mathbb{R}$  be a linear transformation such that

$$T(x^2 + x) = -1; \quad T(x^2 - x) = 1; \quad T(x^2 + 1) = 3.$$

Find  $T(4x^2 + 5x - 3)$ .

## Solution ( first )

Suppose  $a(x^2 + x) + b(x^2 - x) + c(x^2 + 1) = 4x^2 + 5x - 3$ . Then

$$(a + b + c)x^2 + (a - b)x + c = 4x^2 + 5x - 3.$$

## Problem

Let  $T : \mathcal{P}_2 \rightarrow \mathbb{R}$  be a linear transformation such that

$$T(x^2 + x) = -1; \quad T(x^2 - x) = 1; \quad T(x^2 + 1) = 3.$$

Find  $T(4x^2 + 5x - 3)$ .

## Solution ( first )

Suppose  $a(x^2 + x) + b(x^2 - x) + c(x^2 + 1) = 4x^2 + 5x - 3$ . Then

$$(a + b + c)x^2 + (a - b)x + c = 4x^2 + 5x - 3.$$

Solving for  $a$ ,  $b$ , and  $c$  results in the unique solution  $a = 6$ ,  $b = 1$ ,  $c = -3$ .

## Problem

Let  $T : \mathcal{P}_2 \rightarrow \mathbb{R}$  be a linear transformation such that

$$T(x^2 + x) = -1; \quad T(x^2 - x) = 1; \quad T(x^2 + 1) = 3.$$

Find  $T(4x^2 + 5x - 3)$ .

## Solution ( first )

Suppose  $a(x^2 + x) + b(x^2 - x) + c(x^2 + 1) = 4x^2 + 5x - 3$ . Then

$$(a + b + c)x^2 + (a - b)x + c = 4x^2 + 5x - 3.$$

Solving for  $a$ ,  $b$ , and  $c$  results in the unique solution  $a = 6$ ,  $b = 1$ ,  $c = -3$ .  
Thus

$$\begin{aligned} T(4x^2 + 5x - 3) &= T(6(x^2 + x) + (x^2 - x) - 3(x^2 + 1)) \\ &= 6T(x^2 + x) + T(x^2 - x) - 3T(x^2 + 1) \\ &= 6(-1) + 1 - 3(3) = -14. \end{aligned}$$



## Solution ( second )

Notice that  $S = \{x^2 + x, x^2 - x, x^2 + 1\}$  is a basis of  $\mathcal{P}_2$ , and thus  $x^2$ ,  $x$ , and  $1$  can each be written as a linear combination of elements of  $S$ .

## Solution ( second )

Notice that  $S = \{x^2 + x, x^2 - x, x^2 + 1\}$  is a basis of  $\mathcal{P}_2$ , and thus  $x^2$ ,  $x$ , and  $1$  can each be written as a linear combination of elements of  $S$ .

$$x^2 = \frac{1}{2}(x^2 + x) + \frac{1}{2}(x^2 - x)$$

$$x = \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x)$$

$$1 = (x^2 + 1) - \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x).$$

## Solution ( second )

Notice that  $S = \{x^2 + x, x^2 - x, x^2 + 1\}$  is a basis of  $\mathcal{P}_2$ , and thus  $x^2$ ,  $x$ , and  $1$  can each be written as a linear combination of elements of  $S$ .

$$\begin{aligned}x^2 &= \frac{1}{2}(x^2 + x) + \frac{1}{2}(x^2 - x) \\x &= \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x) \\1 &= (x^2 + 1) - \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x).\end{aligned}$$

↓

$$\begin{aligned}\mathbb{T}(x^2) &= \mathbb{T}\left(\frac{1}{2}(x^2 + x) + \frac{1}{2}(x^2 - x)\right) = \frac{1}{2}\mathbb{T}(x^2 + x) + \frac{1}{2}\mathbb{T}(x^2 - x) \\&= \frac{1}{2}(-1) + \frac{1}{2}(1) = 0.\end{aligned}$$

## Solution ( second )

Notice that  $S = \{x^2 + x, x^2 - x, x^2 + 1\}$  is a basis of  $\mathcal{P}_2$ , and thus  $x^2$ ,  $x$ , and  $1$  can each be written as a linear combination of elements of  $S$ .

$$\begin{aligned}x^2 &= \frac{1}{2}(x^2 + x) + \frac{1}{2}(x^2 - x) \\x &= \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x) \\1 &= (x^2 + 1) - \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x).\end{aligned}$$

↓

$$\begin{aligned}\mathbb{T}(x^2) &= \mathbb{T}\left(\frac{1}{2}(x^2 + x) + \frac{1}{2}(x^2 - x)\right) = \frac{1}{2}\mathbb{T}(x^2 + x) + \frac{1}{2}\mathbb{T}(x^2 - x) \\&= \frac{1}{2}(-1) + \frac{1}{2}(1) = 0. \\ \mathbb{T}(x) &= \mathbb{T}\left(\frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x)\right) = \frac{1}{2}\mathbb{T}(x^2 + x) - \frac{1}{2}\mathbb{T}(x^2 - x) \\&= \frac{1}{2}(-1) - \frac{1}{2}(1) = -1.\end{aligned}$$

## Solution ( second )

Notice that  $S = \{x^2 + x, x^2 - x, x^2 + 1\}$  is a basis of  $\mathcal{P}_2$ , and thus  $x^2$ ,  $x$ , and  $1$  can each be written as a linear combination of elements of  $S$ .

$$\begin{aligned}x^2 &= \frac{1}{2}(x^2 + x) + \frac{1}{2}(x^2 - x) \\x &= \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x) \\1 &= (x^2 + 1) - \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x).\end{aligned}$$

↓

$$\begin{aligned}T(x^2) &= T\left(\frac{1}{2}(x^2 + x) + \frac{1}{2}(x^2 - x)\right) = \frac{1}{2}T(x^2 + x) + \frac{1}{2}T(x^2 - x) \\&= \frac{1}{2}(-1) + \frac{1}{2}(1) = 0.\end{aligned}$$

$$\begin{aligned}T(x) &= T\left(\frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x)\right) = \frac{1}{2}T(x^2 + x) - \frac{1}{2}T(x^2 - x) \\&= \frac{1}{2}(-1) - \frac{1}{2}(1) = -1.\end{aligned}$$

$$\begin{aligned}T(1) &= T\left((x^2 + 1) - \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x)\right) \\&= T(x^2 + 1) - \frac{1}{2}T(x^2 + x) - \frac{1}{2}T(x^2 - x) \\&= 3 - \frac{1}{2}(-1) - \frac{1}{2}(1) = 3.\end{aligned}$$



## Solution ( second )

Notice that  $S = \{x^2 + x, x^2 - x, x^2 + 1\}$  is a basis of  $\mathcal{P}_2$ , and thus  $x^2$ ,  $x$ , and  $1$  can each be written as a linear combination of elements of  $S$ .

$$\begin{aligned}x^2 &= \frac{1}{2}(x^2 + x) + \frac{1}{2}(x^2 - x) \\x &= \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x) \\1 &= (x^2 + 1) - \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x).\end{aligned}$$

↓

$$\begin{aligned}T(x^2) &= T\left(\frac{1}{2}(x^2 + x) + \frac{1}{2}(x^2 - x)\right) = \frac{1}{2}T(x^2 + x) + \frac{1}{2}T(x^2 - x) \\&= \frac{1}{2}(-1) + \frac{1}{2}(1) = 0.\end{aligned}$$

$$\begin{aligned}T(x) &= T\left(\frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x)\right) = \frac{1}{2}T(x^2 + x) - \frac{1}{2}T(x^2 - x) \\&= \frac{1}{2}(-1) - \frac{1}{2}(1) = -1.\end{aligned}$$

$$\begin{aligned}T(1) &= T\left((x^2 + 1) - \frac{1}{2}(x^2 + x) - \frac{1}{2}(x^2 - x)\right) \\&= T(x^2 + 1) - \frac{1}{2}T(x^2 + x) - \frac{1}{2}T(x^2 - x) \\&= 3 - \frac{1}{2}(-1) - \frac{1}{2}(1) = 3.\end{aligned}$$

↓

$$T(4x^2 + 5x - 3) = 4T(x^2) + 5T(x) - 3T(1) = 4(0) + 5(-1) - 3(3) = -14.$$



## Remark

The advantage of the second solution over the first is that if you were now asked to find  $T(-6x^2 - 13x + 9)$ , it is easy to use  $T(x^2) = 0$ ,  $T(x) = -1$  and  $T(1) = 3$ :

$$\begin{aligned}T(-6x^2 - 13x + 9) &= -6T(x^2) - 13T(x) + 9T(1) \\ &= -6(0) - 13(-1) + 9(3) = 13 + 27 = 40.\end{aligned}$$

## Remark

The advantage of the second solution over the first is that if you were now asked to find  $T(-6x^2 - 13x + 9)$ , it is easy to use  $T(x^2) = 0$ ,  $T(x) = -1$  and  $T(1) = 3$ :

$$\begin{aligned}T(-6x^2 - 13x + 9) &= -6T(x^2) - 13T(x) + 9T(1) \\ &= -6(0) - 13(-1) + 9(3) = 13 + 27 = 40.\end{aligned}$$

More generally,

$$\begin{aligned}T(ax^2 + bx + c) &= aT(x^2) + bT(x) + cT(1) \\ &= a(0) + b(-1) + c(3) = -b + 3c.\end{aligned}$$

### Definition (Equality of linear transformations)

Let  $V$  and  $W$  be vector spaces, and let  $S$  and  $T$  be linear transformations from  $V$  to  $W$ . Then  $S = T$  if and only if,

$$S(\vec{v}) = T(\vec{v}) \quad \text{for every } \vec{v} \in V.$$

## Definition (Equality of linear transformations)

Let  $V$  and  $W$  be vector spaces, and let  $S$  and  $T$  be linear transformations from  $V$  to  $W$ . Then  $S = T$  if and only if,

$$S(\vec{v}) = T(\vec{v}) \quad \text{for every } \vec{v} \in V.$$

## Theorem

Let  $V$  and  $W$  be vector spaces, where

$$V = \text{span}\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}.$$

Suppose that  $S$  and  $T$  are linear transformations from  $V$  to  $W$ . If  $S(\vec{v}_i) = T(\vec{v}_i)$  for all  $i$ ,  $1 \leq i \leq n$ , then  $S = T$ .

### Definition (Equality of linear transformations)

Let  $V$  and  $W$  be vector spaces, and let  $S$  and  $T$  be linear transformations from  $V$  to  $W$ . Then  $S = T$  if and only if,

$$S(\vec{v}) = T(\vec{v}) \quad \text{for every } \vec{v} \in V.$$

### Theorem

Let  $V$  and  $W$  be vector spaces, where

$$V = \text{span}\{\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n\}.$$

Suppose that  $S$  and  $T$  are linear transformations from  $V$  to  $W$ . If  $S(\vec{v}_i) = T(\vec{v}_i)$  for all  $i$ ,  $1 \leq i \leq n$ , then  $S = T$ .

### Remark

This theorem tells us that a linear transformation is completely determined by its actions on a spanning set.

**Proof.**

We must show that  $S(\vec{v}) = T(\vec{v})$  for each  $\vec{v} \in V$ . Let  $\vec{v} \in V$ . Then (since  $V$  is spanned by  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ ), there exist  $k_1, k_2, \dots, k_n \in \mathbb{R}$  so that

$$\vec{v} = k_1\vec{v}_1 + k_2\vec{v}_2 + \dots + k_n\vec{v}_n.$$

**Proof.**

We must show that  $S(\vec{v}) = T(\vec{v})$  for each  $\vec{v} \in V$ . Let  $\vec{v} \in V$ . Then (since  $V$  is spanned by  $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$ ), there exist  $k_1, k_2, \dots, k_n \in \mathbb{R}$  so that

$$\vec{v} = k_1\vec{v}_1 + k_2\vec{v}_2 + \cdots + k_n\vec{v}_n.$$

It follows that

$$\begin{aligned} S(\vec{v}) &= S(k_1\vec{v}_1 + k_2\vec{v}_2 + \cdots + k_n\vec{v}_n) \\ &= k_1S(\vec{v}_1) + k_2S(\vec{v}_2) + \cdots + k_nS(\vec{v}_n) \\ &= k_1T(\vec{v}_1) + k_2T(\vec{v}_2) + \cdots + k_nT(\vec{v}_n) \\ &= T(k_1\vec{v}_1 + k_2\vec{v}_2 + \cdots + k_n\vec{v}_n) \\ &= T(\vec{v}). \end{aligned}$$

Therefore,  $S = T$ . ■



What is a Linear Transformations


Examples and Problems

Properties of Linear Transformations

**Constructing Linear Transformations**




# Constructing Linear Transformations



## Theorem

Let  $V$  and  $W$  be vector spaces, let  $B = \{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  be a basis of  $V$ , and let  $\vec{w}_1, \vec{w}_2, \dots, \vec{w}_n$  be (not necessarily distinct) vectors of  $W$ .

# Constructing Linear Transformations




## Theorem

Let  $V$  and  $W$  be vector spaces, let  $B = \{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  be a basis of  $V$ , and let  $\vec{w}_1, \vec{w}_2, \dots, \vec{w}_n$  be (not necessarily distinct) vectors of  $W$ . Then

1. There exists a linear transformation  $T : V \rightarrow W$  such that  $T(\vec{b}_i) = \vec{w}_i$  for each  $i$ ,  $1 \leq i \leq n$ ;

# Constructing Linear Transformations




## Theorem

Let  $V$  and  $W$  be vector spaces, let  $B = \{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  be a basis of  $V$ , and let  $\vec{w}_1, \vec{w}_2, \dots, \vec{w}_n$  be (not necessarily distinct) vectors of  $W$ . Then

1. There exists a linear transformation  $T : V \rightarrow W$  such that  $T(\vec{b}_i) = \vec{w}_i$  for each  $i$ ,  $1 \leq i \leq n$ ;
2. This transformation is unique;

# Constructing Linear Transformations



## Theorem

Let  $V$  and  $W$  be vector spaces, let  $B = \{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  be a basis of  $V$ , and let  $\vec{w}_1, \vec{w}_2, \dots, \vec{w}_n$  be (not necessarily distinct) vectors of  $W$ . Then

1. There exists a linear transformation  $T : V \rightarrow W$  such that  $T(\vec{b}_i) = \vec{w}_i$  for each  $i$ ,  $1 \leq i \leq n$ ;
2. This transformation is unique;
3. If

$$\vec{v} = k_1 \vec{b}_1 + k_2 \vec{b}_2 + \dots + k_n \vec{b}_n$$

is a vector of  $V$ , then

$$T(\vec{v}) = k_1 \vec{w}_1 + k_2 \vec{w}_2 + \dots + k_n \vec{w}_n.$$

**Proof.**

Suppose  $\vec{v} \in V$ . Since  $B$  is a basis, there exist unique scalars  $k_1, k_2, \dots, k_n \in \mathbb{R}$  so that  $\vec{v} = k_1\vec{b}_1 + k_2\vec{b}_2 + \dots + k_n\vec{b}_n$ . We now **define**  $T : V \rightarrow W$  by

$$T(\vec{v}) = k_1\vec{w}_1 + k_2\vec{w}_2 + \dots + k_n\vec{w}_n$$

for each  $\vec{v} = k_1\vec{b}_1 + k_2\vec{b}_2 + \dots + k_n\vec{b}_n$  in  $V$ . From this definition,  $T(\vec{b}_i) = \vec{w}_i$  for each  $i$ ,  $1 \leq i \leq n$ .

To prove that  $T$  is a linear transformation, prove that  $T$  preserves addition and scalar multiplication. Let  $\vec{v}, \vec{u} \in V$ . Then

$$\vec{v} = k_1\vec{b}_1 + k_2\vec{b}_2 + \dots + k_n\vec{b}_n \quad \text{and} \quad \vec{u} = l_1\vec{b}_1 + l_2\vec{b}_2 + \dots + l_n\vec{b}_n$$

for some  $k_1, k_2, \dots, k_n \in \mathbb{R}$  and  $l_1, l_2, \dots, l_n \in \mathbb{R}$ .

Proof. (continued)

$$\begin{aligned}T(\vec{v} + \vec{u}) &= T[(k_1\vec{b}_1 + k_2\vec{b}_2 + \cdots + k_n\vec{b}_n) + (\ell_1\vec{b}_1 + \ell_2\vec{b}_2 + \cdots + \ell_n\vec{b}_n)] \\&= T[(k_1 + \ell_1)\vec{b}_1 + (k_2 + \ell_2)\vec{b}_2 + \cdots + (k_n + \ell_n)\vec{b}_n] \\&= (k_1 + \ell_1)\vec{w}_1 + (k_2 + \ell_2)\vec{w}_2 + \cdots + (k_n + \ell_n)\vec{w}_n \\&= (k_1\vec{w}_1 + k_2\vec{w}_2 + \cdots + k_n\vec{w}_n) + (\ell_1\vec{w}_1 + \ell_2\vec{w}_2 + \cdots + \ell_n\vec{w}_n) \\&= T(k_1\vec{b}_1 + k_2\vec{b}_2 + \cdots + k_n\vec{b}_n) + T(\ell_1\vec{b}_1 + \ell_2\vec{b}_2 + \cdots + \ell_n\vec{b}_n) \\&= T(\vec{v}) + T(\vec{u}).\end{aligned}$$

Therefore,  $T$  preserves addition.



Proof. (continued)

$$\begin{aligned}T(\vec{v} + \vec{u}) &= T[(k_1\vec{b}_1 + k_2\vec{b}_2 + \cdots + k_n\vec{b}_n) + (\ell_1\vec{b}_1 + \ell_2\vec{b}_2 + \cdots + \ell_n\vec{b}_n)] \\&= T[(k_1 + \ell_1)\vec{b}_1 + (k_2 + \ell_2)\vec{b}_2 + \cdots + (k_n + \ell_n)\vec{b}_n] \\&= (k_1 + \ell_1)\vec{w}_1 + (k_2 + \ell_2)\vec{w}_2 + \cdots + (k_n + \ell_n)\vec{w}_n \\&= (k_1\vec{w}_1 + k_2\vec{w}_2 + \cdots + k_n\vec{w}_n) + (\ell_1\vec{w}_1 + \ell_2\vec{w}_2 + \cdots + \ell_n\vec{w}_n) \\&= T(k_1\vec{b}_1 + k_2\vec{b}_2 + \cdots + k_n\vec{b}_n) + T(\ell_1\vec{b}_1 + \ell_2\vec{b}_2 + \cdots + \ell_n\vec{b}_n) \\&= T(\vec{v}) + T(\vec{u}).\end{aligned}$$

Therefore,  $T$  preserves addition. Let  $\vec{v}$  be as already defined and let  $r \in \mathbb{R}$ . Then

$$\begin{aligned}T(r\vec{v}) &= T[r(k_1\vec{b}_1 + k_2\vec{b}_2 + \cdots + k_n\vec{b}_n)] \\&= T[(rk_1)\vec{b}_1 + (rk_2)\vec{b}_2 + \cdots + (rk_n)\vec{b}_n] \\&= (rk_1)\vec{w}_1 + (rk_2)\vec{w}_2 + \cdots + (rk_n)\vec{w}_n \\&= r(k_1\vec{w}_1 + k_2\vec{w}_2 + \cdots + k_n\vec{w}_n) \\&= rT(k_1\vec{b}_1 + k_2\vec{b}_2 + \cdots + k_n\vec{b}_n) \\&= rT(\vec{v}).\end{aligned}$$

Therefore,  $T$  preserves scalar multiplication.

Proof. (continued)


Finally, the previous Theorem guarantees that  $T$  is unique: since  $B$  is a basis (and hence a spanning set), the action of  $T$  is completely determined by the fact that  $T(\vec{b}_i) = \vec{w}_i$  for each  $i$ ,  $1 \leq i \leq n$ . This completes the proof of the theorem. ■

Proof. (continued)

Finally, the previous Theorem guarantees that  $T$  is unique: since  $B$  is a basis (and hence a spanning set), the action of  $T$  is completely determined by the fact that  $T(\vec{b}_i) = \vec{w}_i$  for each  $i$ ,  $1 \leq i \leq n$ . This completes the proof of the theorem. ■

## Remark


The significance of this Theorem is that it gives us the ability to define linear transformations between vector spaces, a useful tool in what follows.



### Problem

$B = \{1+x, x+x^2, 1+x^2\}$  is a basis of  $\mathcal{P}_2$ . Let

$$A_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad A_3 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$




## Problem

$B = \{1+x, x+x^2, 1+x^2\}$  is a basis of  $\mathcal{P}_2$ . Let

$$A_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad A_3 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

Find a linear transformation  $T : \mathcal{P}_2 \rightarrow \mathbf{M}_{2,2}$  so the

$$T(1+x) = A_1, \quad T(x+x^2) = A_2, \quad \text{and} \quad T(1+x^2) = A_3,$$



## Problem

$B = \{1+x, x+x^2, 1+x^2\}$  is a basis of  $\mathcal{P}_2$ . Let

$$A_1 = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \quad A_2 = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \quad A_3 = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

Find a linear transformation  $T : \mathcal{P}_2 \rightarrow \mathbf{M}_{2,2}$  so the

$$T(1+x) = A_1, \quad T(x+x^2) = A_2, \quad \text{and} \quad T(1+x^2) = A_3,$$

by specifying  $T(a+bx+cx^2)$  for any  $a+bx+cx^2 \in \mathcal{P}_2$ .

## Solution

Notice that  $(1 + x) + (x + x^2) - (1 + x^2) = 2x$ , and thus

$$x = \frac{1}{2}(1 + x) + \frac{1}{2}(x + x^2) - \frac{1}{2}(1 + x^2),$$

$\Downarrow$

$$\begin{aligned} T(x) &= \frac{1}{2}T(1 + x) + \frac{1}{2}T(x + x^2) - \frac{1}{2}T(1 + x^2) \\ &= \frac{1}{2}A_1 + \frac{1}{2}A_2 - \frac{1}{2}A_3 \\ &= \frac{1}{2} \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \frac{1}{2} \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}. \end{aligned}$$

### Solution (continued)

Next,  $1 = (1 + x) - x$ , so  $T(1) = T(1 + x) - T(x)$ , and thus

$$T(1) = A_1 - \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}.$$



### Solution (continued)

Next,  $1 = (1 + x) - x$ , so  $T(1) = T(1 + x) - T(x)$ , and thus

$$T(1) = A_1 - \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}.$$

Finally,  $x^2 = (x + x^2) - x$ , so  $T(x^2) = T(x + x^2) - T(x)$ , and thus

$$T(x^2) = A_2 - \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix}.$$

### Solution (continued)

Next,  $1 = (1 + x) - x$ , so  $T(1) = T(1 + x) - T(x)$ , and thus

$$T(1) = A_1 - \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}.$$

Finally,  $x^2 = (x + x^2) - x$ , so  $T(x^2) = T(x + x^2) - T(x)$ , and thus

$$T(x^2) = A_2 - \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix}.$$

Therefore,

$$\begin{aligned} T(ax + bx + cx^2) &= aT(1) + bT(x) + cT(x^2) \\ &= \frac{a}{2} \begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix} + \frac{b}{2} \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} + \frac{c}{2} \begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix} \\ &= \frac{1}{2} \begin{bmatrix} a + b - c & -a + b + c \\ -a + b + c & a - b + c \end{bmatrix}. \end{aligned}$$



Solution ( Two – sketch )

Since the set  $\{1 + x, x + x^2, 1 + x^2\}$  is a basis of  $\mathcal{P}_2$ , there exists unique representation:

$$\begin{aligned} a + bx + cx^2 &= l_1(1 + x) + l_2(x + x^2) + l_3(1 + x^2) \\ &= (l_1 + l_3) + (l_1 + l_2)x + (l_2 + l_3)x^2 \end{aligned}$$

↓

$$\begin{cases} l_1 + l_3 = a \\ l_1 + l_2 = b \\ l_2 + l_3 = c \end{cases}$$

↓

$$\begin{cases} l_1 = \frac{1}{2}(a + b - c) \\ l_2 = \frac{1}{2}(-a + b + c) \\ l_3 = \frac{1}{2}(a - b + c) \end{cases}$$

Solution ( Two – sketch )

Since the set  $\{1 + x, x + x^2, 1 + x^2\}$  is a basis of  $\mathcal{P}_2$ , there exists unique representation:

$$\begin{aligned} a + bx + cx^2 &= l_1(1 + x) + l_2(x + x^2) + l_3(1 + x^2) \\ &= (l_1 + l_3) + (l_1 + l_2)x + (l_2 + l_3)x^2 \end{aligned}$$

↓

$$\begin{cases} l_1 + l_3 = a \\ l_1 + l_2 = b \\ l_2 + l_3 = c \end{cases}$$

↓


$$\begin{cases} l_1 = \frac{1}{2}(a + b - c) \\ l_2 = \frac{1}{2}(-a + b + c) \\ l_3 = \frac{1}{2}(a - b + c) \end{cases}$$

## Solution (Two – continued)

Hence,

$$\begin{aligned} & T[a + bx + cx^2] \\ & \parallel \\ & T[l_1(1 + x) + l_2(x + x^2) + l_3(1 + x^2)] \\ & \parallel \\ & l_1T[1 + x] + l_2T[x + x^2] + l_3T[1 + x^2] \\ & \parallel \\ & l_1 \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + l_2 \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + l_3 \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \\ & \parallel \\ & \frac{1}{2}(a + b - c) \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix} + \frac{1}{2}(-a + b + c) \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} + \frac{1}{2}(a - b + c) \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \\ & \parallel \\ & = \frac{1}{2} \begin{bmatrix} a + b - c & -a + b + c \\ -a + b + c & a - b + c \end{bmatrix} \end{aligned}$$






### Problem

Let  $V$  be a vector space, and  $T$  be a linear operator on  $V$ , and  $\mathbf{v}, \mathbf{w} \in V$  such that

$$T(\mathbf{v} + \mathbf{w}) = \mathbf{v} - 2\mathbf{w} \quad \text{and} \quad T(2\mathbf{v} - \mathbf{w}) = 2\mathbf{v}.$$



### Problem

Let  $V$  be a vector space, and  $T$  be a linear operator on  $V$ , and  $\mathbf{v}, \mathbf{w} \in V$  such that

$$T(\mathbf{v} + \mathbf{w}) = \mathbf{v} - 2\mathbf{w} \quad \text{and} \quad T(2\mathbf{v} - \mathbf{w}) = 2\mathbf{v}.$$

Find  $T(\mathbf{v})$  and  $T(\mathbf{w})$ .

## Solution

$$\begin{aligned}T(\mathbf{v}) &= T \left[ \frac{1}{3} ([\mathbf{v} + \mathbf{w}] + [2\mathbf{v} - \mathbf{w}]) \right] \\&= \frac{1}{3} T [\mathbf{v} + \mathbf{w}] + \frac{1}{3} T [2\mathbf{v} - \mathbf{w}] \\&= \frac{1}{3} (\mathbf{v} - 2\mathbf{w}) + \frac{2}{3} \mathbf{v} \\&= \mathbf{v} - \frac{2}{3} \mathbf{w}.\end{aligned}$$

Similarly, as an exercise,  $T(\mathbf{w}) = -\frac{4}{3}\mathbf{w}$ .

