

# Math 221: LINEAR ALGEBRA

## Chapter 7. Linear Transformations

### §7-3. Isomorphisms and Composition

Le Chen<sup>1</sup>

Emory University, 2021 Spring

(last updated on 04/19/2021)



Creative Commons License  
(CC BY-NC-SA)

<sup>1</sup>Slides are adapted from those by Karen Seyffarth from University of Calgary.

What is isomorphism?

Proving vector spaces are isomorphic

Characterizing isomorphisms

Composition of transformations

Inverses

## What is isomorphism?

Proving vector spaces are isomorphic

Characterizing isomorphisms

Composition of transformations

Inverses



# What is an isomorphism?

## Example

$\mathcal{P}_1 = \{ax + b \mid a, b \in \mathbb{R}\}$ , has addition and scalar multiplication defined as follows:

$$\begin{aligned}(a_1x + b_1) + (a_2x + b_2) &= (a_1 + a_2)x + (b_1 + b_2), \\ k(a_1x + b_1) &= (ka_1)x + (kb_1),\end{aligned}$$

for all  $(a_1x + b_1), (a_2x + b_2) \in \mathcal{P}_1$  and  $k \in \mathbb{R}$ .

The role of the variable  $x$  is to distinguish  $a_1$  from  $b_1$ ,  $a_2$  from  $b_2$ ,  $(a_1 + a_2)$  from  $(b_1 + b_2)$ , and  $(ka_1)$  from  $(kb_1)$ .

### Example (continued)

This can be accomplished equally well by using vectors in  $\mathbb{R}^2$ .

$$\mathbb{R}^2 = \left\{ \begin{bmatrix} a \\ b \end{bmatrix} \mid a, b \in \mathbb{R} \right\}$$


where addition and scalar multiplication are defined as follows:

$$\begin{bmatrix} a_1 \\ b_1 \end{bmatrix} + \begin{bmatrix} a_2 \\ b_2 \end{bmatrix} = \begin{bmatrix} a_1 + a_2 \\ b_1 + b_2 \end{bmatrix}, \quad k \begin{bmatrix} a_1 \\ b_1 \end{bmatrix} = \begin{bmatrix} ka_1 \\ kb_1 \end{bmatrix}$$

for all  $\begin{bmatrix} a_1 \\ b_1 \end{bmatrix}, \begin{bmatrix} a_2 \\ b_2 \end{bmatrix} \in \mathbb{R}^2$  and  $k \in \mathbb{R}$ .

## Definition


Let  $V$  and  $W$  be vector spaces, and  $T : V \rightarrow W$  a linear transformation.  $T$  is an **isomorphism** if and only if  $T$  is both one-to-one and onto (i.e.,  $\ker(T) = \{\mathbf{0}\}$  and  $\text{im}(T) = W$ ). If  $T : V \rightarrow W$  is an isomorphism, then the vector spaces  $V$  and  $W$  are said to be **isomorphic**, and we write  $V \cong W$ .



General linear transformation  $T$

## Definition

Let  $V$  and  $W$  be vector spaces, and  $T : V \rightarrow W$  a linear transformation.  $T$  is an **isomorphism** if and only if  $T$  is both one-to-one and onto (i.e.,  $\ker(T) = \{\mathbf{0}\}$  and  $\text{im}(T) = W$ ). If  $T : V \rightarrow W$  is an isomorphism, then the vector spaces  $V$  and  $W$  are said to be **isomorphic**, and we write  $V \cong W$ .



Isomorphism  $T$



## Example

The identity operator on any vector space is an isomorphism.

## Example

The identity operator on any vector space is an isomorphism.

## Example

$T : \mathcal{P}_n \rightarrow \mathbb{R}^{n+1}$  defined by

$$T(a_0 + a_1x + a_2x^2 + \cdots + a_nx^n) = \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}$$

for all  $a_0 + a_1x + a_2x^2 + \cdots + a_nx^n \in \mathcal{P}_n$  is an isomorphism. To verify this, prove that **T is a linear transformation** that is **one-to-one** and **onto**.

What is isomorphism?

**Proving vector spaces are isomorphic**

Characterizing isomorphisms

Composition of transformations

Inverses



## Proving isomorphism of vector spaces

### Problem

Prove that  $\mathbf{M}_{22}$  and  $\mathbb{R}^4$  are isomorphic.

## Proving isomorphism of vector spaces

### Problem

Prove that  $\mathbf{M}_{22}$  and  $\mathbb{R}^4$  are isomorphic.

### Proof.

Let  $T : \mathbf{M}_{22} \rightarrow \mathbb{R}^4$  be defined by

$$T \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \text{ for all } \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbf{M}_{22}.$$

# Proving isomorphism of vector spaces

## Problem

Prove that  $\mathbf{M}_{22}$  and  $\mathbb{R}^4$  are isomorphic.

## Proof.

Let  $T : \mathbf{M}_{22} \rightarrow \mathbb{R}^4$  be defined by

$$T \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} \quad \text{for all } \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbf{M}_{22}.$$

It remains to prove that

1.  $T$  is a linear transformation;
2.  $T$  is one-to-one;
3.  $T$  is onto.

Solution (continued – 1. linear transformation)

Let  $A = \begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix}$ ,  $B = \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} \in \mathbf{M}_{22}$  and let  $k \in \mathbb{R}$ . Then

$$T(A) = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} \quad \text{and} \quad T(B) = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix}.$$



### Solution (continued – 1. linear transformation)

Let  $A = \begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix}$ ,  $B = \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} \in \mathbf{M}_{22}$  and let  $k \in \mathbb{R}$ . Then

$$T(A) = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} \quad \text{and} \quad T(B) = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix}.$$

$$\begin{aligned} & \Downarrow \\ T(A+B) &= T \begin{bmatrix} a_1 + b_1 & a_2 + b_2 \\ a_3 + b_3 & a_4 + b_4 \end{bmatrix} = \begin{bmatrix} a_1 + b_1 \\ a_2 + b_2 \\ a_3 + b_3 \\ a_4 + b_4 \end{bmatrix} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} + \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix} = T(A) + T(B) \end{aligned}$$

Solution (continued – 1. linear transformation)

Let  $A = \begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix}$ ,  $B = \begin{bmatrix} b_1 & b_2 \\ b_3 & b_4 \end{bmatrix} \in \mathbf{M}_{22}$  and let  $k \in \mathbb{R}$ . Then

$$T(A) = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} \quad \text{and} \quad T(B) = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix}.$$

$$T(A+B) = T \begin{bmatrix} a_1 + b_1 & a_2 + b_2 \\ a_3 + b_3 & a_4 + b_4 \end{bmatrix} = \begin{bmatrix} a_1 + b_1 \\ a_2 + b_2 \\ a_3 + b_3 \\ a_4 + b_4 \end{bmatrix} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} + \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ b_4 \end{bmatrix} = T(A) + T(B)$$

$T$  preserves addition.

Solution (continued – 1. linear transformation)

Also

$$T(kA) = T \begin{bmatrix} ka_1 & ka_2 \\ ka_3 & ka_4 \end{bmatrix} = \begin{bmatrix} ka_1 \\ ka_2 \\ ka_3 \\ ka_4 \end{bmatrix} = k \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} = kT(A)$$

Solution (continued – 1. linear transformation)

Also

$$T(kA) = T \begin{bmatrix} ka_1 & ka_2 \\ ka_3 & ka_4 \end{bmatrix} = \begin{bmatrix} ka_1 \\ ka_2 \\ ka_3 \\ ka_4 \end{bmatrix} = k \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} = kT(A)$$

↓

T preserves scalar multiplication.

Since T preserves addition and scalar multiplication, T is a linear transformation.

## Solution (continued – 2. One-to-one)

By definition,

$$\begin{aligned} \ker(T) &= \{A \in \mathbf{M}_{22} \mid T(A) = \mathbf{0}\} \\ &= \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a, b, c, d \in \mathbb{R} \text{ and } \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right\}. \end{aligned}$$

## Solution (continued – 2. One-to-one)

By definition,

$$\begin{aligned}\ker(T) &= \{A \in \mathbf{M}_{22} \mid T(A) = \mathbf{0}\} \\ &= \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a, b, c, d \in \mathbb{R} \text{ and } \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right\}.\end{aligned}$$

If  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \ker T$ , then  $a = b = c = d = 0$ , and thus  $\ker(T) = \{\mathbf{0}_{22}\}$ .

## Solution (continued – 2. One-to-one)

By definition,

$$\begin{aligned}\ker(T) &= \{A \in \mathbf{M}_{22} \mid T(A) = \mathbf{0}\} \\ &= \left\{ \begin{bmatrix} a & b \\ c & d \end{bmatrix} \mid a, b, c, d \in \mathbb{R} \text{ and } \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} \right\}.\end{aligned}$$

If  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \ker T$ , then  $a = b = c = d = 0$ , and thus  $\ker(T) = \{\mathbf{0}_{22}\}$ .

↓

T is one-to-one.

Solution (continued – 3. Onto)

Let

$$X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \in \mathbb{R}^4,$$

and define matrix  $A \in \mathbf{M}_{22}$  as follows:

$$A = \begin{bmatrix} x_1 & x_2 \\ x_3 & x_4 \end{bmatrix}.$$



Solution (continued – 3. Onto)

Let

$$X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \in \mathbb{R}^4,$$

and define matrix  $A \in \mathbf{M}_{22}$  as follows:

$$A = \begin{bmatrix} x_1 & x_2 \\ x_3 & x_4 \end{bmatrix}.$$

Then  $T(A) = X$ , and therefore  $T$  is onto.

Finally, since  $T$  is a linear transformation that is one-to-one and onto,  $T$  is an isomorphism.

Solution (continued – 3. Onto)

Let

$$X = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} \in \mathbb{R}^4,$$

and define matrix  $A \in \mathbf{M}_{22}$  as follows:

$$A = \begin{bmatrix} x_1 & x_2 \\ x_3 & x_4 \end{bmatrix}.$$

Then  $T(A) = X$ , and therefore  $T$  is onto.

Finally, since  $T$  is a linear transformation that is one-to-one and onto,  $T$  is an isomorphism. Therefore,  $\mathbf{M}_{22}$  and  $\mathbb{R}^4$  are isomorphic vector spaces. ■

### Example ( Other isomorphic vector spaces )

1. For all integers  $n \geq 0$ ,  $\mathcal{P}_n \cong \mathbb{R}^{n+1}$ .
2. For all integers  $m$  and  $n$ ,  $m, n \geq 1$ ,  $\mathbf{M}_{mn} \cong \mathbb{R}^{m \times n}$ .
3. For all integers  $m$  and  $n$ ,  $m, n \geq 1$ ,  $\mathbf{M}_{mn} \cong \mathcal{P}_{mn-1}$ .

You should be able to define appropriate linear transformations and prove each of these statements.

What is isomorphism?

Proving vector spaces are isomorphic

**Characterizing isomorphisms**

Composition of transformations

Inverses

## Characterizing isomorphisms

# Characterizing isomorphisms

## Theorem

Let  $V$  and  $W$  be finite dimensional vector spaces and  $T : V \rightarrow W$  a linear transformation. The following are equivalent.

1.  $T$  is an isomorphism.
2. If  $\{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  is any basis of  $V$ , then  $\{T(\vec{b}_1), T(\vec{b}_2), \dots, T(\vec{b}_n)\}$  is a basis of  $W$ .
3. There exists a basis  $\{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  of  $V$  such that  $\{T(\vec{b}_1), T(\vec{b}_2), \dots, T(\vec{b}_n)\}$  is a basis of  $W$ .

# Characterizing isomorphisms

## Theorem

Let  $V$  and  $W$  be finite dimensional vector spaces and  $T : V \rightarrow W$  a linear transformation. The following are equivalent.

1.  $T$  is an isomorphism.
2. If  $\{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  is any basis of  $V$ , then  $\{T(\vec{b}_1), T(\vec{b}_2), \dots, T(\vec{b}_n)\}$  is a basis of  $W$ .
3. There exists a basis  $\{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  of  $V$  such that  $\{T(\vec{b}_1), T(\vec{b}_2), \dots, T(\vec{b}_n)\}$  is a basis of  $W$ .

## Proof.

(1)  $\Rightarrow$  (2): This is because

- One-to-one linear transformations preserve independent sets.
- Onto linear transformations preserve spanning sets.

(2)  $\Rightarrow$  (3) is trivial.

Proof. (Continued)

(3)  $\Rightarrow$  (1). We need to prove that  $T$  is both onto and one-to-one.

If  $T(\vec{v}) = \vec{0}$ , write  $\vec{v} = v_1\vec{b}_1 + \cdots + v_n\vec{b}_n$  where each  $v_i$  is in  $\mathbb{R}$ . Then

$$\vec{0} = T(\vec{v}) = v_1T(\vec{b}_1) + \cdots + v_nT(\vec{b}_n)$$

so  $v_1 = \cdots = v_n = 0$  by (3). Hence  $\vec{v} = \vec{0}$ , so  $\ker T = \{\vec{0}\}$  and  $T$  is one-to-one.

To show that  $T$  is onto, let  $\vec{w}$  be any vector in  $W$ . By (3) there exist  $w_1, \dots, w_n$  in  $\mathbb{R}$  such that

$$\vec{w} = w_1T(\vec{b}_1) + \cdots + w_nT(\vec{b}_n) = T(w_1\vec{b}_1 + \cdots + w_n\vec{b}_n)$$

Thus  $T$  is onto. ■



Suppose  $V$  and  $W$  are finite dimensional vector spaces with  $\dim(V) = \dim(W)$ , and let

$$\{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\} \quad \text{and} \quad \{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_n\}$$

be bases of  $V$  and  $W$  respectively.

Suppose  $V$  and  $W$  are finite dimensional vector spaces with  $\dim(V) = \dim(W)$ , and let

$$\{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\} \quad \text{and} \quad \{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_n\}$$

be bases of  $V$  and  $W$  respectively. Then  $T : V \rightarrow W$  defined by

$$T(\vec{b}_i) = \vec{f}_i \text{ for } 1 \leq i \leq n$$

is a **linear transformation** that maps a basis of  $V$  to a basis of  $W$ . By the previous Theorem,  $T$  is an isomorphism.

Suppose  $V$  and  $W$  are finite dimensional vector spaces with  $\dim(V) = \dim(W)$ , and let

$$\{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\} \quad \text{and} \quad \{\vec{f}_1, \vec{f}_2, \dots, \vec{f}_n\}$$

be bases of  $V$  and  $W$  respectively. Then  $T : V \rightarrow W$  defined by

$$T(\vec{b}_i) = \vec{f}_i \text{ for } 1 \leq i \leq n$$

is a **linear transformation** that maps a basis of  $V$  to a basis of  $W$ . By the previous Theorem,  $T$  is an isomorphism.

Conversely, if  $V$  and  $W$  are isomorphic and  $T : V \rightarrow W$  is an isomorphism, then (by the previous Theorem) for any basis  $\{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  of  $V$ ,  $\{T(\vec{b}_1), T(\vec{b}_2), \dots, T(\vec{b}_n)\}$  is a basis of  $W$ , implying that  $\dim(V) = \dim(W)$ .

This proves the next theorem.

## Theorem

Finite dimensional vector spaces  $V$  and  $W$  are isomorphic if and only if  $\dim(V) = \dim(W)$ .

## Theorem

Finite dimensional vector spaces  $V$  and  $W$  are isomorphic if and only if  $\dim(V) = \dim(W)$ .

## Corollary

If  $V$  is a vector space with  $\dim(V) = n$ , then  $V$  is isomorphic to  $\mathbb{R}^n$ .

## Problem

Let  $V$  denote the set of  $2 \times 2$  real symmetric matrices. Then  $V$  is a vector space with dimension three. Find an isomorphism  $T : \mathcal{P}_2 \rightarrow V$  with the property that  $T(1) = I_2$  (the  $2 \times 2$  identity matrix).

## Problem

Let  $V$  denote the set of  $2 \times 2$  real symmetric matrices. Then  $V$  is a vector space with dimension three. Find an isomorphism  $T : \mathcal{P}_2 \rightarrow V$  with the property that  $T(1) = I_2$  (the  $2 \times 2$  identity matrix).

## Solution

$$V = \left\{ \begin{bmatrix} a & b \\ b & c \end{bmatrix} \mid a, b, c \in \mathbb{R} \right\} = \text{span} \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}.$$

## Problem

Let  $V$  denote the set of  $2 \times 2$  real symmetric matrices. Then  $V$  is a vector space with dimension three. Find an isomorphism  $T : \mathcal{P}_2 \rightarrow V$  with the property that  $T(1) = I_2$  (the  $2 \times 2$  identity matrix).

## Solution

$$V = \left\{ \begin{bmatrix} a & b \\ b & c \end{bmatrix} \mid a, b, c \in \mathbb{R} \right\} = \text{span} \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}.$$

Let

$$B = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}.$$

Then  $B$  is independent, and  $\text{span}(B) = V$ , so  $B$  is a basis of  $V$ . Also,  $\dim(V) = 3 = \dim(\mathcal{P}_2)$ .



## Problem

Let  $V$  denote the set of  $2 \times 2$  real symmetric matrices. Then  $V$  is a vector space with dimension three. Find an isomorphism  $T : \mathcal{P}_2 \rightarrow V$  with the property that  $T(1) = I_2$  (the  $2 \times 2$  identity matrix).

## Solution

$$V = \left\{ \begin{bmatrix} a & b \\ b & c \end{bmatrix} \mid a, b, c \in \mathbb{R} \right\} = \text{span} \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}.$$

Let

$$B = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}.$$

Then  $B$  is independent, and  $\text{span}(B) = V$ , so  $B$  is a basis of  $V$ . Also,  $\dim(V) = 3 = \dim(\mathcal{P}_2)$ . However, we want a basis of  $V$  that contains  $I_2$ .

Solution (continued)

Let

$$B' = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}.$$

Since  $B'$  consists of  $\dim(V)$  symmetric independent matrices,  $B'$  is a basis of  $V$ . Note that  $I_2 \in B'$ .

## Solution (continued)

Let

$$B' = \left\{ \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right\}.$$

Since  $B'$  consists of  $\dim(V)$  symmetric independent matrices,  $B'$  is a basis of  $V$ . Note that  $I_2 \in B'$ . Define

$$T(1) = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, T(x) = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, T(x^2) = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}.$$

Then for all  $ax^2 + bx + c \in \mathcal{P}_2$ ,

$$T(ax^2 + bx + c) = \begin{bmatrix} c & b \\ b & a + c \end{bmatrix},$$

and  $T(1) = I_2$ .

By the previous Theorem,  $T : \mathcal{P}_2 \rightarrow V$  is an isomorphism. ■

## Theorem

Let  $V$  and  $W$  be vector spaces, and  $T : V \rightarrow W$  a linear transformation. If  $\dim(V) = \dim(W) = n$ , then  $T$  is an isomorphism if and only if  $T$  is either one-to-one or onto.

## Theorem

Let  $V$  and  $W$  be vector spaces, and  $T : V \rightarrow W$  a linear transformation. If  $\dim(V) = \dim(W) = n$ , then  $T$  is an isomorphism if and only if  $T$  is either one-to-one or onto.

## Proof.

( $\Rightarrow$ ) By definition, an isomorphism is both one-to-one and onto.

## Theorem

Let  $V$  and  $W$  be vector spaces, and  $T : V \rightarrow W$  a linear transformation. If  $\dim(V) = \dim(W) = n$ , then  $T$  is an isomorphism if and only if  $T$  is either one-to-one or onto.

## Proof.

( $\Rightarrow$ ) By definition, an isomorphism is both one-to-one and onto.

( $\Leftarrow$ ) Suppose that  $T$  is one-to-one. Then  $\ker(T) = \{\vec{0}\}$ , so  $\dim(\ker(T)) = 0$ .  
By the Dimension Theorem,

$$\begin{aligned}\dim(V) &= \dim(\operatorname{im}(T)) + \dim(\ker(T)) \\ n &= \dim(\operatorname{im}(T)) + 0\end{aligned}$$

so  $\dim(\operatorname{im}(T)) = n = \dim(W)$ . Furthermore  $\operatorname{im}(T) \subseteq W$ , so it follows that  $\operatorname{im}(T) = W$ . Therefore,  $T$  is onto, and hence is an isomorphism.

Proof. (continued)

( $\Leftarrow$ ) Suppose that  $T$  is onto. Then  $\text{im}(T) = W$ , so  $\dim(\text{im}(T)) = \dim(W) = n$ . By the Dimension Theorem,

$$\begin{aligned}\dim(V) &= \dim(\text{im}(T)) + \dim(\ker(T)) \\ n &= n + \dim(\ker(T))\end{aligned}$$

so  $\dim(\ker(T)) = 0$ . The only vector space with dimension zero is the zero vector space, and thus  $\ker(T) = \{\vec{0}\}$ . Therefore,  $T$  is one-to-one, and hence is an isomorphism. ■

What is isomorphism?

Proving vector spaces are isomorphic

Characterizing isomorphisms

**Composition of transformations**

Inverses





# Composition of transformations

## Definition


Let  $V, W$  and  $U$  be vector spaces, and let

$$T : V \rightarrow W \quad \text{and} \quad S : W \rightarrow U$$

be linear transformations. The **composite** of  $T$  and  $S$  is

$$ST : V \rightarrow U$$

where  $(ST)(\vec{v}) = S(T(\vec{v}))$  for all  $\vec{v} \in V$ . The process of obtaining  $ST$  from  $S$  and  $T$  is called **composition**.



## Example

Let  $S : \mathbf{M}_{22} \rightarrow \mathbf{M}_{22}$  and  $T : \mathbf{M}_{22} \rightarrow \mathbf{M}_{22}$  be linear transformations such that

$$S(A) = -A^T \quad \text{and} \quad T \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{bmatrix} b & a \\ d & c \end{bmatrix} \quad \text{for all } A = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathbf{M}_{22}.$$

Then

$$(ST) \begin{bmatrix} a & b \\ c & d \end{bmatrix} = S \begin{bmatrix} b & a \\ d & c \end{bmatrix} = \begin{bmatrix} -b & -d \\ -a & -c \end{bmatrix},$$

and

$$(TS) \begin{bmatrix} a & b \\ c & d \end{bmatrix} = T \begin{bmatrix} -a & -c \\ -b & -d \end{bmatrix} = \begin{bmatrix} -c & -a \\ -d & -b \end{bmatrix}.$$

If  $a, b, c$  and  $d$  are distinct, then  $(ST)(A) \neq (TS)(A)$ .

This illustrates that, in general,  $ST \neq TS$ .

## Theorem

Let  $V, W, U$  and  $Z$  be vector spaces and

$$V \xrightarrow{T} W \xrightarrow{S} U \xrightarrow{R} Z$$

be linear transformations. Then

1.  $ST$  is a linear transformation.
2.  $T1_V = T$  and  $1_W T = T$ .
3.  $(RS)T = R(ST)$ .

Problem ( The composition of onto transformations is onto )

Let  $V, W$  and  $U$  be vector spaces, and let

$$V \xrightarrow{T} W \xrightarrow{S} U$$

be linear transformations. Prove that if  $T$  and  $S$  are onto, then  $ST$  is onto.

Problem ( The composition of onto transformations is onto )

Let  $V, W$  and  $U$  be vector spaces, and let


$$V \xrightarrow{T} W \xrightarrow{S} U$$

be linear transformations. Prove that if  $T$  and  $S$  are onto, then  $ST$  is onto.

Proof.

Let  $\mathbf{z} \in U$ . Since  $S$  is onto, there exists a vector  $\mathbf{y} \in W$  such that  $S(\mathbf{y}) = \mathbf{z}$ . Furthermore, since  $T$  is onto, there exists a vector  $\mathbf{x} \in V$  such that  $T(\mathbf{x}) = \mathbf{y}$ . Thus

$$\mathbf{z} = S(\mathbf{y}) = S(T(\mathbf{x})) = (ST)(\mathbf{x}),$$

showing that for each  $\mathbf{z} \in U$  there exists and  $\mathbf{x} \in V$  such that  $(ST)(\mathbf{x}) = \mathbf{z}$ . Therefore,  $ST$  is onto. 

Problem ( The composition of one-to-one transformations is one-to-one )

Let  $V, W$  and  $U$  be vector spaces, and let

$$V \xrightarrow{T} W \xrightarrow{S} U$$

be linear transformations. Prove that if  $T$  and  $S$  are one-to-one, then  $ST$  is one-to-one.

Problem ( The composition of one-to-one transformations is one-to-one )

Let  $V, W$  and  $U$  be vector spaces, and let

$$V \xrightarrow{T} W \xrightarrow{S} U$$

be linear transformations. Prove that if  $T$  and  $S$  are one-to-one, then  $ST$  is one-to-one.

The proof of this is left as an exercise.



What is isomorphism?

Proving vector spaces are isomorphic

Characterizing isomorphisms

Composition of transformations

**Inverses**



# Inverses

## Theorem

Let  $V$  and  $W$  be finite dimensional vector spaces, and  $T : V \rightarrow W$  a linear transformation. Then the following statements are equivalent.

1.  $T$  is an isomorphism.
2. There exists a linear transformation  $S : W \rightarrow V$  so that

$$ST = 1_V \quad \text{and} \quad TS = 1_W.$$

In this case, the isomorphism  $S$  is uniquely determined by  $T$ :

$$\text{if } \vec{w} \in W \quad \text{and} \quad \vec{w} = T(\vec{v}), \text{ then } S(\vec{w}) = \vec{v}.$$

# Inverses

## Theorem

Let  $V$  and  $W$  be finite dimensional vector spaces, and  $T : V \rightarrow W$  a linear transformation. Then the following statements are equivalent.

1.  $T$  is an isomorphism.
2. There exists a linear transformation  $S : W \rightarrow V$  so that

$$ST = 1_V \quad \text{and} \quad TS = 1_W.$$

In this case, the isomorphism  $S$  is uniquely determined by  $T$ :

$$\text{if } \vec{w} \in W \quad \text{and} \quad \vec{w} = T(\vec{v}), \text{ then } S(\vec{w}) = \vec{v}.$$

Given an isomorphism  $T : V \rightarrow W$ , the unique isomorphism satisfying the second condition of the theorem is the **inverse** of  $T$ , and is written  $T^{-1}$ .

Remark ( Fundamental Identities (relating  $T$  and  $T^{-1}$ ) )

If  $V$  and  $W$  are vector spaces and  $T : V \rightarrow W$  is an isomorphism, then  $T^{-1} : W \rightarrow V$  is a linear transformation such that

$$(T^{-1}T)(\vec{v}) = \vec{v} \quad \text{and} \quad (TT^{-1})(\vec{w}) = \vec{w}$$

for each  $\vec{v} \in V$ ,  $\vec{w} \in W$ . Equivalently,

$$T^{-1}T = 1_V \quad \text{and} \quad TT^{-1} = 1_W.$$

## Problem

The function  $T : \mathcal{P}_2 \rightarrow \mathbb{R}^3$  defined by

$$T(a + bx + cx^2) = \begin{bmatrix} a - c \\ 2b \\ a + c \end{bmatrix} \text{ for all } a + bx + cx^2 \in \mathcal{P}_2$$

is a linear transformation (this is left for you to verify). Does  $T$  have an inverse? If so, find  $T^{-1}$ .

## Solution

Since  $\dim(\mathcal{P}_2) = 3 = \dim(\mathbb{R}^3)$ , it suffices to prove that  $T$  is either one-to-one or onto.

## Solution

Since  $\dim(\mathcal{P}_2) = 3 = \dim(\mathbb{R}^3)$ , it suffices to prove that  $T$  is either one-to-one or onto.

Suppose  $a + bx + cx^2 \in \ker(T)$ . Then

$$\begin{cases} a - c = 0 \\ 2b = 0 \\ a + c = 0 \end{cases} \implies \begin{cases} a = 0 \\ b = 0 \\ c = 0 \end{cases}$$



## Solution

Since  $\dim(\mathcal{P}_2) = 3 = \dim(\mathbb{R}^3)$ , it suffices to prove that  $T$  is either one-to-one or onto.

Suppose  $a + bx + cx^2 \in \ker(T)$ . Then

$$\begin{cases} a - c = 0 \\ 2b = 0 \\ a + c = 0 \end{cases} \implies \begin{cases} a = 0 \\ b = 0 \\ c = 0 \end{cases}$$

Therefore,  $\ker(T) = \{\mathbf{0}\}$ , and hence  $T$  is one-to-one. By our earlier observation, it follows that  $T$  is onto, and thus is an isomorphism.

### Solution (continued)

To find  $T^{-1}$ , we need to specify  $T^{-1} \begin{bmatrix} p \\ q \\ r \end{bmatrix}$  for any  $\begin{bmatrix} p \\ q \\ r \end{bmatrix} \in \mathbb{R}^3$ .

Let  $a + bx + cx^2 \in \mathcal{P}_2$ , and suppose

$$T(a + bx + cx^2) = \begin{bmatrix} p \\ q \\ r \end{bmatrix}.$$

By the definition of  $T$ ,  $p = a - c$ ,  $q = 2b$  and  $r = a + c$ . We now solve for  $a$ ,  $b$  and  $c$  in terms of  $p$ ,  $q$  and  $r$ .

$$\left[ \begin{array}{ccc|c} 1 & 0 & -1 & p \\ 0 & 2 & 0 & q \\ 1 & 0 & 1 & r \end{array} \right] \rightarrow \cdots \rightarrow \left[ \begin{array}{ccc|c} 1 & 0 & 0 & (r+p)/2 \\ 0 & 1 & 0 & q/2 \\ 0 & 0 & 1 & (r-p)/2 \end{array} \right].$$

### Solution (continued)

We now have  $a = \frac{r+p}{2}$ ,  $b = \frac{q}{2}$  and  $c = \frac{r-p}{2}$ , and thus

$$T(a + bx + cx^2) = \begin{bmatrix} p \\ q \\ r \end{bmatrix} = T\left(\frac{r+p}{2} + \frac{q}{2}x + \frac{r-p}{2}x^2\right)$$

## Solution (continued)

We now have  $a = \frac{r+p}{2}$ ,  $b = \frac{q}{2}$  and  $c = \frac{r-p}{2}$ , and thus

$$T(a + bx + cx^2) = \begin{bmatrix} p \\ q \\ r \end{bmatrix} = T\left(\frac{r+p}{2} + \frac{q}{2}x + \frac{r-p}{2}x^2\right)$$

Therefore,

$$\begin{aligned} T^{-1} \begin{bmatrix} p \\ q \\ r \end{bmatrix} &= T^{-1}\left(T\left(\frac{r+p}{2} + \frac{q}{2}x + \frac{r-p}{2}x^2\right)\right) \\ &= (T^{-1}T)\left(\frac{r+p}{2} + \frac{q}{2}x + \frac{r-p}{2}x^2\right) \\ &= \frac{r+p}{2} + \frac{q}{2}x + \frac{r-p}{2}x^2. \end{aligned}$$



## Definition

Let  $V$  be a vector space with  $\dim(V) = n$ , let  $B = \{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  be a fixed basis of  $V$ , and let  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  denote the standard basis of  $\mathbb{R}^n$ .

## Definition

Let  $V$  be a vector space with  $\dim(V) = n$ , let  $B = \{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  be a fixed basis of  $V$ , and let  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  denote the standard basis of  $\mathbb{R}^n$ . We define a transformation  $C_B : V \rightarrow \mathbb{R}^n$  by

$$C_B(a_1\vec{b}_1 + a_2\vec{b}_2 + \cdots + a_n\vec{b}_n) = a_1\vec{e}_1 + a_2\vec{e}_2 + \cdots + a_n\vec{e}_n = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}.$$

## Definition

Let  $V$  be a vector space with  $\dim(V) = n$ , let  $B = \{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  be a fixed basis of  $V$ , and let  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  denote the standard basis of  $\mathbb{R}^n$ . We define a transformation  $C_B : V \rightarrow \mathbb{R}^n$  by

$$C_B(a_1\vec{b}_1 + a_2\vec{b}_2 + \dots + a_n\vec{b}_n) = a_1\vec{e}_1 + a_2\vec{e}_2 + \dots + a_n\vec{e}_n = \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix}.$$

Then  $C_B$  is a linear transformation such that  $C_B(\vec{b}_i) = \vec{e}_i$ ,  $1 \leq i \leq n$ , and thus  $C_B$  is an isomorphism, called **the coordinate isomorphism corresponding to  $B$** .

## Example

Let  $V$  be a vector space and let  $B = \{\vec{b}_1, \vec{b}_2, \dots, \vec{b}_n\}$  be a fixed basis of  $V$ . Then  $C_B : V \rightarrow \mathbb{R}^n$  is invertible, and it is clear that  $C_B^{-1} : \mathbb{R}^n \rightarrow V$  is defined by

$$C_B^{-1} \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} = a_1 \vec{b}_1 + a_2 \vec{b}_2 + \cdots + a_n \vec{b}_n \text{ for each } \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} \in \mathbb{R}^n.$$