

# Math 362: Mathematical Statistics II

Le Chen

le.chen@emory.edu

Emory University  
Atlanta, GA

Last updated on April 13, 2021

2021 Spring

# Chapter 7. Inference Based on The Normal Distribution

§ 7.1 Introduction

§ 7.2 Comparing  $\frac{\bar{Y}-\mu}{\sigma/\sqrt{n}}$  and  $\frac{\bar{Y}-\mu}{S/\sqrt{n}}$

§ 7.3 Deriving the Distribution of  $\frac{\bar{Y}-\mu}{S/\sqrt{n}}$

§ 7.4 Drawing Inferences About  $\mu$

§ 7.5 Drawing Inferences About  $\sigma^2$

# Chapter 7. Inference Based on The Normal Distribution

§ 7.1 Introduction

§ 7.2 Comparing  $\frac{\bar{Y}-\mu}{\sigma/\sqrt{n}}$  and  $\frac{\bar{Y}-\mu}{S/\sqrt{n}}$

§ 7.3 Deriving the Distribution of  $\frac{\bar{Y}-\mu}{S/\sqrt{n}}$

§ 7.4 Drawing Inferences About  $\mu$

§ 7.5 Drawing Inferences About  $\sigma^2$

For a random sample of size  $n$  from  $N(\mu, \sigma^2)$ :

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i - \bar{Y})^2$$

$\Downarrow$

$$\frac{(n-1)S^2}{\sigma^2} = \frac{1}{\sigma^2} \sum_{i=1}^n (Y_i - \bar{Y})^2 \sim \text{Chi Square}(n-1)$$

$$\mathbb{P}\left(\chi_{\alpha/2, n-1}^2 \leq \frac{(n-1)S^2}{\sigma^2} \leq \chi_{1-\alpha/2, n-1}^2\right) = 1 - \alpha.$$

100(1 -  $\alpha$ )% C.I. for  $\sigma^2$ :

$$\left( \frac{(n-1)S^2}{\chi_{1-\alpha/2, n-1}^2}, \frac{(n-1)S^2}{\chi_{\alpha/2, n-1}^2} \right)$$

100(1 -  $\alpha$ )% C.I. for  $\sigma$ :

$$\left( \sqrt{\frac{(n-1)S^2}{\chi_{1-\alpha/2, n-1}^2}}, \sqrt{\frac{(n-1)S^2}{\chi_{\alpha/2, n-1}^2}} \right)$$

Testing  $H_0 : \sigma^2 = \sigma_0^2$

v.s.

(at the  $\alpha$  level of significance)

$$\chi^2 = \frac{(n-1)s^2}{\sigma_0^2}$$

$H_1 : \sigma^2 < \sigma_0^2:$

Reject  $H_0$  if

$$\chi^2 \leq \chi_{\alpha, n-1}^2$$

$H_1 : \sigma^2 \neq \sigma_0^2:$

Reject  $H_0$  if

$$\chi^2 \leq \chi_{\alpha/2, n-1}^2 \text{ or}$$

$$\chi^2 \geq \chi_{1-\alpha/2, n-1}^2$$

$H_1 : \sigma^2 > \sigma_0^2:$

Reject  $H_0$  if

$$\chi^2 \geq \chi_{1-\alpha, n-1}^2$$

**E.g. 1.** The width of a confidence interval for  $\sigma^2$  is a function of  $n$  and  $S^2$ :

$$W = \frac{(n-1)S^2}{\chi_{\alpha/2, n-1}^2} - \frac{(n-1)S^2}{\chi_{1-\alpha/2, n-1}^2}$$

Find the smallest  $n$  such that the average width of a 95% C.I. for  $\sigma^2$  is no greater than  $0.8\sigma^2$ .

**Sol.** Notice that  $\mathbb{E}[S^2] = \sigma^2$ . Hence, we need to find  $n$  s.t.

$$(n-1) \left( \frac{1}{\chi_{0.025, n-1}^2} - \frac{1}{\chi_{0.975, n-1}^2} \right) \leq 0.8.$$

Trial and error (numerics on R) gives  $n = 57$ .

```

1 > # Example 7.5.1
2 > n=seq(45,60,1)
3 > l=qchisq(0.025,n-1)
4 > u=qchisq(0.975,n-1)
5 > e=(n-1)* (1/l-1/u)
6 > m=cbind(n,l,u,e)
7 > colnames(m) = c("n",
8 +               "chi(0.025,n-1)",
9 +               "chi(0.975,n-1)",
10 +              "error")
11 > m
12      n chi(0.025,n-1) chi(0.975,n-1) error
13 [1,] 45      27.57457      64.20146 0.9103307
14 [2,] 46      28.36615      65.41016 0.8984312
15 [3,] 47      29.16005      66.61653 0.8869812
16 [4,] 48      29.95620      67.82065 0.8759533
17 [5,] 49      30.75451      69.02259 0.8653224
18 [6,] 50      31.55492      70.22241 0.8550654
19 [7,] 51      32.35736      71.42020 0.8451612
20 [8,] 52      33.16179      72.61599 0.8355901
21 [9,] 53      33.96813      73.80986 0.8263340
22 [10,] 54     34.77633      75.00186 0.8173761
23 [11,] 55     35.58634      76.19205 0.8087008
24 [12,] 56     36.39811      77.38047 0.8002937
25
26 [13,] 57     37.21159      78.56716 0.7921414
27 [14,] 58     38.02674      79.75219 0.7842313
28 [15,] 59     38.84351      80.93559 0.7765517
29 [16,] 60     39.66186      82.11741 0.7690918

```

## Case Study 7.5.2

Mutual funds are investment vehicles consisting of a portfolio of various types of investments. If such an investment is to meet annual spending needs, the owner of shares in the fund is interested in the average of the annual returns of the fund. Investors are also concerned with the volatility of the annual returns, measured by the variance or standard deviation. One common method of evaluating a mutual fund is to compare it to a benchmark, the Lipper Average being one of these. This index number is the average of returns from a universe of mutual funds.

The Global Rock Fund is a typical mutual fund, with heavy investments in international funds. It claimed to best the Lipper Average in terms of volatility over the period from 1989 through 2007. Its returns are given in the table below.

Year	Investment Return %	Year	Investment Return %
1989	15.32	1999	27.43
1990	1.62	2000	8.57
1991	28.43	2001	1.88
1992	11.91	2002	-7.96
1993	20.71	2003	35.98
1994	-2.15	2004	14.27
1995	23.29	2005	10.33
1996	15.96	2006	15.94
1997	11.12	2007	16.71
1998	0.37		

The standard deviation for these returns is 11.28%, while the corresponding figure for the Lipper Average is 11.67%. Now, clearly, the Global Rock Fund has a smaller standard deviation than the Lipper Average, but is this small difference due just to random variation? The hypothesis test is meant to answer such questions.



$$H_0 : \sigma^2 = (11.67)^2$$

versus

$$H_1 : \sigma^2 < (11.67)^2$$

Let  $\alpha = 0.05$ . With  $n = 19$ , the critical value for the chi square ratio [from part (b) of Theorem 7.5.2] is  $\chi_{1-\alpha, n-1}^2 = \chi_{0.95, 18}^2 = 9.390$  (see Figure 7.5.3). But

$$\chi^2 = \frac{(n-1)s^2}{\sigma_0^2} = \frac{(19-1)(11.28)^2}{(11.67)^2} = 16.82$$

so our decision is clear: Do not reject  $H_0$ .

