

DISCRETE MATHEMATICS
SEMINAR

Off-diagonal Ramsey numbers for slowly growing hypergraphs

Jiaxi Nie, PhD
Georgia Tech

Abstract: For a k -uniform hypergraph F and a positive integer n , the Ramsey number $r(F, n)$ denotes the minimum N such that every N -vertex F -free k -uniform hypergraph contains an independent set of n vertices. A hypergraph is *slowly growing* if there is an ordering e_1, e_2, \dots, e_t of its edges such that $|e_i \setminus \bigcup_{j=1}^{i-1} e_j| \leq 1$ for each $i \in \{2, \dots, t\}$. We prove that if $k \geq 3$ is fixed and F is any non k -partite slowly growing k -uniform hypergraph, then for $n \geq 2$,

$$r(F, n) = \Omega\left(\frac{n^k}{(\log n)^{2k-2}}\right).$$

In particular, we deduce that the off-diagonal Ramsey number $r(F_5, n)$ is of order $n^3/\text{polylog}(n)$, where F_5 is the triple system $\{123, 124, 345\}$. This is the only 3-uniform Berge triangle for which the polynomial power of its off-diagonal Ramsey number was not previously known. Our constructions use pseudorandom graphs, martingales, and hypergraph containers.

Thursday, October 31, 2024, 4:00 pm
Mathematics and Science Center: MSC W303

MATHEMATICS
EMORY UNIVERSITY